
antibodies - online.com







anti-WTAP antibody (C-Term)



Image



()	ve	K\ /		A .
	\cup	1 V/	-	V۷

0.00000		
Quantity:	ty: 400 µL	
Target:	WTAP	
Binding Specificity:	AA 272-301, C-Term	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This WTAP antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
lmmunogen:	This WTAP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 272-301 amino acids from the C-terminal region of human WTAP.	
Clone:	RB30382	
Isotype:	lg Fraction	
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.	
Target Details		
Target:	WTAP	
Alternative Name:	WTAP (WTAP Products)	
Background:	The Wilms tumor suppressor gene WT1 appears to play a role in both transcriptional and	

Target Details

posttranscriptional regulation of certain cellular genes. This gene encodes a WT1-associating
protein, which is a ubiquitously expressed nuclear protein. Like WT1 protein, this protein is
localized throughout the nucleoplasm as well as in speckles and partially colocalizes with
splicing factors. Alternative splicing of this gene results in three transcript variants, two of
which encode the same isoform.

Molecular Weight:	44244
Gene ID:	9589
NCBI Accession:	NP_001257460, NP_001257461, NP_001257462, NP_004897, NP_690596, NP_690597


UniProt: Q15007

Application Details

Application Notes:	WB: 1:1000
Restrictions:	For Research Use only

Handling

Format:	Liquid	
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	WTAP Antibody (C-term) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, keep at -20 °C.	
Expiry Date:	6 months	



Western Blotting

Image 1. WT Antibody (C-term) 16885b western blot analysis in Jurkat cell line lysates (35 μ g/lane).This demonstrates the WT antibody detected the WT protein (arrow).