



[Go to Product page](#)

Datasheet for ABIN1537121  
**anti-XKRX antibody (C-Term)**

1 Image

Overview

Quantity:	400 µL
Target:	XKRX
Binding Specificity:	AA 317-345, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This XKRX antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	This XKRX antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 317-345 amino acids from the C-terminal region of human XKRX.
Clone:	RB35922
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	XKRX
Alternative Name:	XKRX ( <a href="#">XKRX Products</a> )
Background:	This gene encodes a protein that is related to a component of the XK/Kell complex of the Kell

## Target Details

---

blood group system. The encoded protein includes several transmembrane domains, is known to be exposed to the cell surface, and may function as a membrane transporter.

Molecular Weight: 52052

Gene ID: 402415

NCBI Accession: [NP\\_997724](#)

UniProt: [Q6PP77](#)

## Application Details

---

Application Notes: WB: 1:1000

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: XKRX Antibody (C-term) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, keep at -20 °C.

Expiry Date: 6 months



### Western Blotting

**Image 1.** All lanes : Anti-XKRX Antibody (C-term) at 1:1000 dilution Lane 1: human fetal kidney lysate Lane 2: NCI-whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 52 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.