

Datasheet for ABIN1537263
anti-ONECUT3 antibody (C-Term)[Go to Product page](#)

1 Image

Overview

| | |
|----------------------|--|
| Quantity: | 400 µL |
| Target: | ONECUT3 |
| Binding Specificity: | AA 393-419, C-Term |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This ONECUT3 antibody is un-conjugated |
| Application: | Western Blotting (WB) |

Product Details

| | |
|-----------------------|--|
| Immunogen: | This ONECUT3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 393-419 amino acids from the C-terminal region of human ONECUT3. |
| Clone: | RB37735 |
| Isotype: | Ig Fraction |
| Predicted Reactivity: | M |
| Purification: | This antibody is purified through a protein A column, followed by peptide affinity purification. |

Target Details

| | |
|-------------------|--|
| Target: | ONECUT3 |
| Alternative Name: | ONECUT3 (ONECUT3 Products) |

Target Details


| | |
|-------------------|---|
| Background: | ONECUT3 is transcriptional activator. It binds the consensus DNA sequence 5'-DHWATTGAYTWW-3' on a variety of gene promoters such as those of HNF3B and TTR (By similarity). |
| Molecular Weight: | 50037 |
| Gene ID: | 390874 |
| NCBI Accession: | NP_001073957 |
| UniProt: | O60422 |

Application Details

| | |
|--------------------|-----------------------|
| Application Notes: | WB: 1:1000 |
| Restrictions: | For Research Use only |

Handling

| | |
|--------------------|--|
| Format: | Liquid |
| Buffer: | Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C, -20 °C |
| Storage Comment: | ONECUT3 Antibody (C-term) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, keep at -20 °C. |
| Expiry Date: | 6 months |



Western Blotting

Image 1. Anti-ONECUT3 Antibody (C-term) at 1:1000 dilution + MDA-MB-231 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 50 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.