



[Go to Product page](#)

Datasheet for ABIN1537747

## anti-STAG3L1 antibody (AA 108-136)

### 2 Images

#### Overview

Quantity:	400 µL
Target:	STAG3L1
Binding Specificity:	AA 108-136
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This STAG3L1 antibody is un-conjugated
Application:	Western Blotting (WB)

#### Product Details

Immunogen:	This STAG3L1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 108-136 amino acids from the Central region of human STAG3L1.
Clone:	RB39888
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Target Details

Target:	STAG3L1
Alternative Name:	STAG3L1 ( <a href="#">STAG3L1 Products</a> )
Background:	The function of this protein remains unknown.

## Target Details

Molecular Weight: 23775

Gene ID: 54441

UniProt: [P0CL83](#)

## Application Details

Application Notes: WB: 1:2000. WB: 1:2000

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: STAG3L1 Antibody (Center) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, keep at -20 °C.


Expiry Date: 6 months

## Images



### Western Blotting

**Image 1.** All lanes : Anti-STAG3L1 Antibody (Center) at 1:2000 dilution Lane 1: MCF-7 whole cell lysate Lane 2: ZR-75-1 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 24 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.



### Western Blotting

**Image 2.** All lanes : Anti-STAG3L1 Antibody (Center) at 1:2000 dilution Lane 1: human testis lysate Lane 2: MCF-7 whole cell lysate Lane 3: MOLT-4 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 24 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.