
antibodies - online.com







anti-RNF128 antibody (AA 115-143)



Image



\sim	
()\/\	rview
\cup	

Quantity:	400 μL
Target:	RNF128
Binding Specificity:	AA 115-143
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RNF128 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	This RNF128 antibody is generated from rabbits immunized with a KLH conjugated synthetic
	peptide between 115-143 amino acids from the Central region of human RNF128.
Clone:	RB40302
Isotype:	Ig Fraction
Predicted Reactivity:	В
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target:	RNF128

Target Details

Background:

The protein encoded by this gene is a type I transmembrane protein that localizes to the endocytic pathway. This protein contains a RING zinc-finger motif and has been shown to possess E3 ubiquitin ligase activity. Expression of this gene in retrovirally transduced T cell hybridoma significantly inhibits activation-induced IL2 and IL4 cytokine production. Induced expression of this gene was observed in anergic CD4(+) T cells, which suggested a role in the induction of anergic phenotype. Alternatively spliced transcript variants encoding distinct isoforms have been reported.

Molecular Weight:

46521

Gene ID:

79589

NCBI Accession:

NP_078815, NP_919445

UniProt:

Q8TEB7

Application Details

Application Notes:

WB: 1:1000

Restrictions:

For Research Use only

Handling

F	0	r	r	n	а	t	

Liquid

Buffer:

Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative:

Sodium azide

Precaution of Use:

This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

should be handled by trained staff only.

Storage:

4 °C,-20 °C


Storage Comment:

RNF128 Antibody (Center) can be refrigerated at 2-8 °C for up to 6 months. For long term

storage, keep at -20 °C.

Expiry Date:

6 months



Western Blotting

Image 1. RN Antibody (Center) (ABIN1537774 and ABIN2838343) western blot analysis in MDA-M cell line lysates (35 μ g/lane). This demonstrates the RN antibody detected the RN protein (arrow).