

Datasheet for ABIN1538175
anti-RNF133 antibody (AA 245-271)[Go to Product page](#)

2 Images

Overview

| | |
|----------------------|---------------------------------------|
| Quantity: | 400 µL |
| Target: | RNF133 |
| Binding Specificity: | AA 245-271 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This RNF133 antibody is un-conjugated |
| Application: | Western Blotting (WB) |

Product Details

| | |
|---------------|---|
| Immunogen: | This RNF133 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 245-271 amino acids from the Central region of human RNF133. |
| Clone: | RB37620 |
| Isotype: | Ig Fraction |
| Purification: | This antibody is purified through a protein A column, followed by peptide affinity purification. |

Target Details

| | |
|-------------------|---|
| Target: | RNF133 |
| Alternative Name: | RNF133 (RNF133 Products) |
| Background: | The protein encoded by this gene contains a RING finger domain, a motif present in a variety of |

Target Details


| | |
|-------------------|---|
| | functionally distinct proteins and known to be involved in protein-protein and protein-DNA interactions. This gene has no intron. [provided by RefSeq]. |
| Molecular Weight: | 42294 |
| Gene ID: | 168433 |
| NCBI Accession: | NP_631914 |
| UniProt: | Q8WVZ7 |

Application Details

| | |
|--------------------|------------------------|
| Application Notes: | WB: 1:1000. WB: 1:1000 |
| Restrictions: | For Research Use only |


Handling

| | |
|--------------------|--|
| Format: | Liquid |
| Buffer: | Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | RNF133 Antibody (Center) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, keep at -20 °C. |
| Expiry Date: | 6 months |



Western Blotting

Image 1. Anti-RN Antibody (Center) at 1:1000 dilution + K562 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 42 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.



Western Blotting

Image 2. RN Antibody (Center) (ABIN1538175 and ABIN2849189) western blot analysis in NCI- cell line lysates (35 µg/lane). This demonstrates the RN antibody detected the RN protein (arrow).