# antibodies - online.com







# anti-PCDHGB4 antibody (AA 353-379)



Image



$\sim$				
	$ V \cap$	r\/I	19	٨

Quantity:	400 μL
Target:	PCDHGB4
Binding Specificity:	AA 353-379
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PCDHGB4 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	This PCDHGB4 antibody is generated from rabbits immunized with a KLH conjugated synthetic
Immunogen:	This PCDHGB4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 353-379 amino acids from the Central region of human PCDHGB4.
Immunogen: Clone:	
	peptide between 353-379 amino acids from the Central region of human PCDHGB4.
Clone:	peptide between 353-379 amino acids from the Central region of human PCDHGB4.  RB39608
Clone:  Isotype:  Purification:	peptide between 353-379 amino acids from the Central region of human PCDHGB4.  RB39608  Ig Fraction
Clone:  Isotype:  Purification:  Target Details	peptide between 353-379 amino acids from the Central region of human PCDHGB4.  RB39608  Ig Fraction  This antibody is purified through a protein A column, followed by peptide affinity purification.
Clone:  Isotype:  Purification:	peptide between 353-379 amino acids from the Central region of human PCDHGB4.  RB39608  Ig Fraction
Clone: Isotype: Purification: Target Details	peptide between 353-379 amino acids from the Central region of human PCDHGB4.  RB39608  Ig Fraction  This antibody is purified through a protein A column, followed by peptide affinity purification.

tandemly linked on chromosome five. These gene clusters have an immunoglobulin-like organization, suggesting that a novel mechanism may be involved in their regulation and expression. The gamma gene cluster includes 22 genes divided into 3 subfamilies. Subfamily A contains 12 genes, subfamily B contains 7 genes and 2 pseudogenes, and the more distantly related subfamily C contains 3 genes. The tandem array of 22 large, variable region exons are followed by a constant region, containing 3 exons shared by all genes in the cluster. Each variable region exon encodes the extracellular region, which includes 6 cadherin ectodomains and a transmembrane region. The constant region exons encode the common cytoplasmic region. These neural cadherin-like cell adhesion proteins most likely play a critical role in the establishment and function of specific cell-cell connections in the brain. This particular family member is expressed in fibroblasts and is thought to play a role in wound healing in response to injury. Alternative splicing has been described for the gamma cluster genes.


Molecular Weight:	99927
Gene ID:	8641
NCBI Accession:	NP_003727, NP_115269
UniProt:	Q9UN71

#### **Application Details**

Application Notes:	WB: 1:1000
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	PCDHGB4 Antibody (Center) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, keep at -20 °C.
Expiry Date:	6 months



## **Western Blotting**

**Image 1.** PCDHGB4 Antibody (Center) (ABIN1538347 and ABIN2850269) western blot analysis in Uterus tissue lysates (35  $\mu$ g/lane).This demonstrates the PCDHGB4 antibody detected the PCDHGB4 protein (arrow).