# antibodies .- online.com







# anti-Calicin antibody (AA 373-400)



Image



#### Overview

Quantity:	400 μL
Target:	Calicin (CCIN)
Binding Specificity:	AA 373-400
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Calicin antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	This CCIN antibody is generated from rabbits immunized with a KLH conjugated synthetic
	peptide between 373-400 amino acids from the Central region of human CCIN.
Clone:	RB37671
Isotype:	Ig Fraction
Predicted Reactivity:	B, M, Rat
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Target Details	
Target Details  Target:	Calicin (CCIN)

## **Target Details**


Background:	The protein encoded by this gene is a basic protein of the sperm head cytoskeleton. This protein contains kelch repeats and a BTB/POZ domain and is necessary for normal morphology during sperm differentiation. This gene is intronless.
Gene ID:	881
NCBI Accession:	NP_005884
UniProt:	Q13939

## Application Details

Application Notes:	WB: 1:1000
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	CCIN Antibody (Center) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, keep at -20 °C.
Expiry Date:	6 months



#### **Western Blotting**

Image 1. All lanes: Anti-CCIN Antibody (Center) at 1:1000 dilution Lane 1: human testis lysate Lane 2: HL-60 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 67 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.