

Datasheet for ABIN1539148  
**anti-Multicilin antibody (N-Term)**[Go to Product page](#)

## 1 Image

## Overview

Quantity:	400 µL
Target:	Multicilin (MCIDAS)
Binding Specificity:	AA 20-46, N-Term
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Multicilin antibody is un-conjugated
Application:	Western Blotting (WB)

## Product Details

Immunogen:	This IDAS antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 20-46 amino acids from the N-terminal region of human IDAS.
Clone:	RB30498
Isotype:	Ig Fraction
Predicted Reactivity:	B
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

## Target Details

Target:	Multicilin (MCIDAS)
Alternative Name:	IDAS ( <a href="#">MCIDAS Products</a> )

## Target Details


Background:	Transcription regulator required for multiciliate cell differentiation. Acts by promoting transcription of genes required for multiciliate cell formation. Probably acts in a multiprotein complex By similarity. Plays a role in mitotic cell cycle progression by promoting cell cycle exit. Ref.1
Molecular Weight:	41720
Gene ID:	345643
NCBI Accession:	<a href="#">NP_001177716</a>
UniProt:	<a href="#">D6RGH6</a>

## Application Details

Application Notes:	WB: 1:1000
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	IDAS Antibody (N-term) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, keep at -20 °C.
Expiry Date:	6 months



Western Blotting

**Image 1.** All lanes : Anti-IDAS Antibody (N-term) at 1:1000 dilution Lane 1: U-20S whole cell lysate Lane 2: T47D whole cell lysate Lane 3: HeLa whole cell lysate Lane 4: Rat liver whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 42 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.