antibodies -online.com





anti-WBP2NL antibody (N-Term)



Image



()	11/0	r\ /1	$\triangle 1 $
	$\lor \lor \vdash$	$I \vee I$	ew

Quantity:	400 μL
Target:	WBP2NL
Binding Specificity:	AA 9-36, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB)
Product Details	

Immunogen:	This WBP2NL antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 9-36 amino acids from the N-terminal region of human WBP2NL.
Clone:	RB37771
Isotype:	lg Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	WBP2NL	
Alternative Name:	WBP2NL (WBP2NL Products)	
Background: WBP2NL is a sperm-specific WW domain-binding protein that promotes meiotic read and pronuclear development during oocyte fertilization (Wu et al., 2007 [PubMed		
	17289678]).[supplied by OMIM].	

Target Details

Molecular Weight:	31909
Gene ID:	164684
NCBI Accession:	NP_689826
UniProt:	Q6ICG8


Application Details

Application Notes:	WB: 1:1000
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	WBP2NL Antibody (N-term) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, keep at -20 °C.
Expiry Date:	6 months

Images



Western Blotting

Image 1. Anti-WBP2NL Antibody (N-term) at 1:1000 dilution + human testis lysate Lysates/proteins at 20 μg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 32 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.