

Datasheet for ABIN1539751 anti-KIT antibody (pTyr730)

1 Image



Quantity:

Overview

400 µL

Target:

KIT

Binding Specificity:

pTyr730

Reactivity:

Human

Host:

Rabbit

Polyclonal

Clonality:
Conjugate:

This KIT antibody is un-conjugated

Application:

Dot Blot (DB)

Product Details

Immunogen:

This KIT Antibody is generated from rabbits immunized with a KLH conjugated synthetic

phosphopeptide corresponding to amino acid residues surrounding Y730 of human KIT.

Clone:

RB39719

Isotype:

Ig Fraction

Predicted Reactivity:

B, Pig

Purification:

This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:


KIT

Alternative Name:

KIT (KIT Products)

Target Details

This gene encodes the human homolog of the proto-oncogene c-kit. C-kit was first identified as the cellular homolog of the feline sarcoma viral oncogene v-kit. This protein is a type 3 transmembrane receptor for MGF (mast cell growth factor, also known as stem cell factor). Mutations in this gene are associated with gastrointestinal stromal tumors, mast cell disease, acute myelogenous lukemia, and piebaldism. Multiple transcript variants encoding different isoforms have been found for this gene.
109865
3815
NP_000213, NP_001087241
P10721
RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Sensory Perception of Sound, Stem Cell Maintenance, Production of Molecular Mediator of Immune Response, Regulation of long-term Neuronal Synaptic Plasticity
DB: 1:500
For Research Use only
Liquid
Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Sodium azide
This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
4 °C,-20 °C
Phospho-KIT-Y730 Antibody can be refrigerated at 2-8 °C for up to 6 months. For long term storage, keep at -20 °C.
6 months



Dot Blot

Image 1. Dot blot analysis of Phospho-KIT- Antibody Phospho-specific Pab m on nitrocellulose membrane. 50 ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.6 μ g per ml.