

Datasheet for ABIN1568566

beta-2 Microglobulin ELISA Kit





Overview

in mouse serum, plasma, urine, tissue homogenates, cell lysates, cell culture supernates an other biological fluids. Sample Type: Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Homogenate, Urine Analytical Method: Quantitative Detection Method: Colorimetric Specificity: This assay has high sensitivity and excellent specificity for detection of this index. Cross-Reactivity (Details): No significant cross-reactivity or interference between this index and analogues was observed. Note: Limited by current skills and knowledge, it is impossible for us to complete the cross-		
Reactivity: Mouse Method Type: Sandwich ELISA Detection Range: 0.156 ng/mL - 10 ng/mL Minimum Detection Limit: 0.156 ng/mL Application: ELISA Product Details Purpose: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of by in mouse serum, plasma, urine, tissue homogenates, cell lysates, cell culture supernates an other biological fluids. Sample Type: Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Homogenate, Urine Analytical Method: Quantitative Detection Method: Colorimetric Specificity: This assay has high sensitivity and excellent specificity for detection of this index. Cross-Reactivity (Details): No significant cross-reactivity or interference between this index and analogues was observated by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between this index and all the analogues, therefore, cross reaction may exist.	Quantity:	96 tests
Method Type: Sandwich ELISA Detection Range: 0.156 ng/mL - 10 ng/mL Minimum Detection Limit: 0.156 ng/mL Application: ELISA Product Details Purpose: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of bin in mouse serum, plasma, urine, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Sample Type: Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Homogenate, Urine Analytical Method: Quantitative Detection Method: Colorimetric Specificity: This assay has high sensitivity and excellent specificity for detection of this index. Cross-Reactivity (Details): No significant cross-reactivity or interference between this index and analogues was observing the conservative of the cross-reactivity detection between this index and all the analogues, therefore, cross reaction may exist.	Target:	beta-2 Microglobulin (B2M)
Detection Range: 0.156 ng/mL - 10 ng/mL Minimum Detection Limit: 0.156 ng/mL Application: ELISA Product Details The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of being in mouse serum, plasma, urine, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Sample Type: Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Homogenate, Urine Analytical Method: Quantitative Detection Method: Colorimetric Specificity: This assay has high sensitivity and excellent specificity for detection of this index. No significant cross-reactivity or interference between this index and analogues was obsert Note: Limited by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between this index and all the analogues, therefore, cross reaction may exist.	Reactivity:	Mouse
Minimum Detection Limit: 0.156 ng/mL Application: ELISA Product Details Purpose: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of bin mouse serum, plasma, urine, tissue homogenates, cell lysates, cell culture supernates an other biological fluids. Sample Type: Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Homogenate, Urine Analytical Method: Quantitative Detection Method: Colorimetric Specificity: This assay has high sensitivity and excellent specificity for detection of this index. Cross-Reactivity (Details): No significant cross-reactivity or interference between this index and analogues was obsert Note: Limited by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between this index and all the analogues, therefore, cross reaction may exist.	Method Type:	Sandwich ELISA
Application: Product Details Purpose: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of bid in mouse serum, plasma, urine, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Sample Type: Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Homogenate, Urine Analytical Method: Quantitative Detection Method: Colorimetric Specificity: This assay has high sensitivity and excellent specificity for detection of this index. Cross-Reactivity (Details): No significant cross-reactivity or interference between this index and analogues was obsert Note: Limited by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between this index and all the analogues, therefore, cross reaction may exist.	Detection Range:	0.156 ng/mL - 10 ng/mL
Product Details Purpose: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of by in mouse serum, plasma, urine, tissue homogenates, cell lysates, cell culture supernates an other biological fluids. Sample Type: Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Homogenate, Urine Analytical Method: Quantitative Detection Method: Colorimetric Specificity: This assay has high sensitivity and excellent specificity for detection of this index. Cross-Reactivity (Details): No significant cross-reactivity or interference between this index and analogues was obsernated by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between this index and all the analogues, therefore, cross reaction may exist.	Minimum Detection Limit:	0.156 ng/mL
Purpose: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of but in mouse serum, plasma, urine, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Sample Type: Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Homogenate, Urine Analytical Method: Quantitative Detection Method: Colorimetric Specificity: This assay has high sensitivity and excellent specificity for detection of this index. Cross-Reactivity (Details): No significant cross-reactivity or interference between this index and analogues was observance. Note: Limited by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between this index and all the analogues, therefore, cross reaction may exist.	Application:	ELISA
in mouse serum, plasma, urine, tissue homogenates, cell lysates, cell culture supernates an other biological fluids. Sample Type: Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Homogenate, Urine Analytical Method: Quantitative Detection Method: Colorimetric Specificity: This assay has high sensitivity and excellent specificity for detection of this index. Cross-Reactivity (Details): No significant cross-reactivity or interference between this index and analogues was observance. Note: Limited by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between this index and all the analogues, therefore, cross reaction may exist.	Product Details	
Analytical Method: Detection Method: Colorimetric Specificity: This assay has high sensitivity and excellent specificity for detection of this index. Cross-Reactivity (Details): No significant cross-reactivity or interference between this index and analogues was observed. Note: Limited by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between this index and all the analogues, therefore, cross reaction may exist.	Purpose:	The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of b2M in mouse serum, plasma, urine, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Detection Method: Colorimetric This assay has high sensitivity and excellent specificity for detection of this index. Cross-Reactivity (Details): No significant cross-reactivity or interference between this index and analogues was observed. Note: Limited by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between this index and all the analogues, therefore, cross reaction may exist.	Sample Type:	Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Homogenate, Urine
Specificity: This assay has high sensitivity and excellent specificity for detection of this index. Cross-Reactivity (Details): No significant cross-reactivity or interference between this index and analogues was observed. Note: Limited by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between this index and all the analogues, therefore, cross reaction may exist.	Analytical Method:	Quantitative
Cross-Reactivity (Details): No significant cross-reactivity or interference between this index and analogues was observed. Note: Limited by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between this index and all the analogues, therefore, cross reaction may exist.	Detection Method:	Colorimetric
Note: Limited by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between this index and all the analogues, therefore, cross reaction may exist.	Specificity:	This assay has high sensitivity and excellent specificity for detection of this index.
Sensitivity: 0.052 ng/mL	Cross-Reactivity (Details):	No significant cross-reactivity or interference between this index and analogues was observed. Note: Limited by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between this index and all the analogues, therefore, cross reaction may still exist.
	Sensitivity:	0.052 ng/mL

Product Details

Components:

- · Pre-coated, ready to use 96-well strip plate
- · Standard (freeze dried)
- · Standard Diluent
- · Detection Reagent A
- · Detection Reagent B
- · Assay Diluent A
- · Assay Diluent B
- TMB
- · Stop Solution
- Wash Buffer (30X)
- Plate sealer for 96 wells
- · Instruction manual

Material not included:

- 1. Microplate reader with 450 \pm 10nm filter.
- 2. Precision single or multi-channel pipettes and disposable tips.
- 3. Eppendorf Tubes for diluting samples.
- 4. Deionized or distilled water.
- 5. Absorbent paper for blotting the microtiter plate.
- 6. Container for Wash Solution.

Target Details

Target:	beta-2 Microglobulin (B2M)
Alternative Name:	b2M (B2M Products)
Background:	Alternative name: BMG, B2MG
Gene ID:	12010
UniProt:	P01887
Pathways:	TCR Signaling, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune
	Effector Process

Application Details

Sample Volume:	100 μL
Assay Time:	1 - 4.5 h
Plate:	Pre-coated Pre-coated
Protocol:	1. Prepare all reagents, samples and standards 2. Add 100µL standard or sample to each well. Incubate 2 hours at 37°C

	3. Aspirate and add 100µL prepared Detection Reagent A. Incubate 1 hour at 37°C
	4. Aspirate and wash 3 times
	5. Add 100µL prepared Detection Reagent B. Incubate 1 hour at 37°C
	6. Aspirate and wash 5 times
	7. Add 90µL Substrate Solution. Incubate 15-25 minutes at 37°C
	8. Add 50µL Stop Solution. Read at 450nm immediately.
Assay Procedure:	The microtiter plate provided in this kit has been pre-coated with an antibody specific to the
	index. Standards or samples are then added to the appropriate microtiter plate wells with a
	biotin-conjugated antibody preparation specific to the index. Next, Avidin conjugated to
	Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB
	substrate solution is added, only those wells that contain the index, biotin-conjugated antibody
	and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is
	terminated by the addition of sulphuric acid solution and the color change is measured
	spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of the index in
	the samples is then determined by comparing the O.D. of the samples to the standard curve.
Assay Precision:	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level
	the index were tested 20 times on one plate, respectively.
	 Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level the index were tested on 3 different plates, 8 replicates in each plate.
	• CV(%) = SD/meanX100
	Intra-assay: CV<10%
	Inter-assay: CV<12%
Restrictions:	For Research Use only
Handling	
Precaution of Use:	The Stop Solution suggested for use with this kit is an acid solution. Wear eye, hand, face, and
	clothing protection when using this material.
Handling Advice:	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less
	than 5 % within the expiration date under appropriate storage conditions. Note: To minimize
	unnecessary influences on the performance, operation procedures and lab conditions,
	especially room temperature, air humidity and incubator temperatures should be strictly
	regulated. It is also strongly suggested that the whole assay is performed by the same
	experimenter from the beginning to the end.

Handling


Storage Comment:

The Assay Plate, Standard, Detection Reagent A and Detection Reagent B should be stored at -20°C upon being received. After receiving the kit, Substrate should be always stored at 4°C.Other reagents are kept according to the labels on vials. But for long term storage, please keep the whole kit at -20°C. The unused strips should be kept in a sealed bag with the desiccant provided to minimize exposure to damp air. The test kit may be used throughout the expiration date of the kit (six months from the date of manufacture). Opened test kits will remain stable until the expiring date shown, provided it is stored as prescribed above.

Expiry Date:

12 months

Images



ELISA

Image 1. Typical Standard Curve