

Datasheet for ABIN1571557

NPSR1 ELISA Kit

1 Image



Go to Product page

	ve	rvi	0	W
\cup	VC	I V I	\sim	v v

Quantity:	96 tests	
Target:	NPSR1	
Reactivity:	Human	
Method Type:	Sandwich ELISA	
Detection Range:	0.312 ng/mL - 20 ng/mL	
Minimum Detection Limit:	0.312 ng/mL	
Application:	ELISA	
Product Details		
Purpose:	The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of NPSR in human tissue homogenates and other biological fluids.	
Sample Type:	Tissue Homogenate	
Analytical Method:	Quantitative	
Detection Method:	Colorimetric	
Specificity:	This assay has high sensitivity and excellent specificity for detection of this index.	
Cross-Reactivity (Details):	No significant cross-reactivity or interference between this index and analogues was observed. Note: Limited by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between this index and all the analogues, therefore, cross reaction may still exist.	
Sensitivity:	0.108 ng/mL	

Product Details

Components:

- Pre-coated, ready to use 96-well strip plate
- · Standard (freeze dried)
- · Standard Diluent
- · Detection Reagent A
- · Detection Reagent B
- · Assay Diluent A
- · Assay Diluent B
- TMB
- · Stop Solution
- Wash Buffer (30X)
- · Plate sealer for 96 wells
- Instruction manual

Material not included:

- 1. Microplate reader with 450 \pm 10nm filter.
- 2. Precision single or multi-channel pipettes and disposable tips.
- 3. Eppendorf Tubes for diluting samples.
- 4. Deionized or distilled water.
- 5. Absorbent paper for blotting the microtiter plate.
- 6. Container for Wash Solution.

Target Details

Target:	NPSR1	
Alternative Name:	NPSR (NPSR1 Products)	
Background:	Alternative name: ASRT2, GPR154, GPRA, NPSR1, PGR14, VRR1, G Protein-Coupled Receptor For Asthma Susceptibility, G Protein-Coupled Receptor 154, G-protein coupled receptor PGR14	
Gene ID:	387129	
UniProt:	Q6W5P4	

Application Details		
Sample Volume:	100 μL	
Assay Time:	1 - 4.5 h	
Plate:	Pre-coated	
Protocol:	1. Prepare all reagents, samples and standards 2. Add 100µL standard or sample to each well. Incubate 2 hours at 37°C	
	3. Aspirate and add 100µL prepared Detection Reagent A. Incubate 1 hour at 37°C	
	4. Aspirate and wash 3 times	

	5. Add 100µL prepared Detection Reagent B. Incubate 1 hour at 37°C		
	6. Aspirate and wash 5 times		
	7. Add 90µL Substrate Solution. Incubate 15-25 minutes at 37°C		
	8. Add 50µL Stop Solution. Read at 450nm immediately.		
Assay Procedure:	The microtiter plate provided in this kit has been pre-coated with an antibody specific to the		
	index. Standards or samples are then added to the appropriate microtiter plate wells with a		
	biotin-conjugated antibody preparation specific to the index. Next, Avidin conjugated to		
	Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB		
	substrate solution is added, only those wells that contain the index, biotin-conjugated antibody		
	and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is		
	terminated by the addition of sulphuric acid solution and the color change is measured		
	spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of the index in		
	the samples is then determined by comparing the O.D. of the samples to the standard curve.		
Assay Precision:	 Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level the index were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level the index were tested on 3 different plates. 8 replicates in each plate. 		
	 the index were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100 		
	• Intra-assay: CV<10%		
	Inter-assay: CV<12%		
Restrictions:	For Research Use only		
Handling			
Precaution of Use:	The Stop Solution suggested for use with this kit is an acid solution. Wear eye, hand, face, and		
	clothing protection when using this material.		
Handling Advice:	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less		
	than 5 % within the expiration date under appropriate storage conditions. Note: To minimize		
	unnecessary influences on the performance, operation procedures and lab conditions,		
	especially room temperature, air humidity and incubator temperatures should be strictly		
	regulated. It is also strongly suggested that the whole assay is performed by the same		
	experimenter from the beginning to the end.		
Storage:	4 °C,-20 °C		
Ctorogo Commont:	The Assay Plate, Standard, Detection Reagent A and Detection Reagent B should be stored at -		
Storage Comment:	The Assay Hate, Standard, Detection Neagent A and Detection Neagent D should be stored at		


Handling

4°C.Other reagents are kept according to the labels on vials. But for long term storage, please keep the whole kit at -20°C. The unused strips should be kept in a sealed bag with the desiccant provided to minimize exposure to damp air. The test kit may be used throughout the expiration date of the kit (six months from the date of manufacture). Opened test kits will remain stable until the expiring date shown, provided it is stored as prescribed above.

Expiry Date:

12 months

Images



ELISA

Image 1.