

[Go to Product page](#)

Datasheet for ABIN1589610

FLT1 Protein (glycosylated, Monomer, Soluble)

Overview

Quantity:	5 µg
Target:	FLT1
Protein Characteristics:	glycosylated, Monomer, Soluble
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Biological Activity:	Active

Product Details

Purpose:	VEGFR-1/Flt-1 (D5), soluble
Sequence:	SKLKDPELSL KGTQHIMQAG QTLHLQCRGE AAHKWSLPEM VSKESERLSI TKSACGRNGK QFCSTLTLNT AQANHTGFYS CKYLAVPTSK KKETESAIYI FISDTGRPFV EMYSEIPEII HMTEGRELVI PCRVTSPNIT VTLKKFPLDT LIPDGKRIIW DSRKGFISN ATYKEIGLLT CEATVNGHLY KTNYLTHRQT NTIIDVQIST PRPVKLLRGH TLVLNCTATT PLNTRVQMTW SYPDEKNKRA SVRRRIDQSN SHANIFYSVL TIDKMQNKDK GLYTICRVRSG PSFKSVNTSV HIYDKAFITV KHRKQQVLET VAGKRSYRLS MKVKAFPSPE VVWLKDGLPA TEKSARYLTR GYSLIKDVT EEDAGNYTIL LSIKQSNVFK NLTATLIVNV KPQIYEKAVS SFPDPALYPL GSRQILTCTA YGIPQPTIKW FWHPCNHNHS EARCDFCSNN EESFILDADS NMGNRIESIT QRMAIIEGKN KMASTLVVAD SRISGIYICI ASNKVGTVGR NISFYITDVP NGFHVN
Specificity:	Chromosomal location:13q12
Characteristics:	Length (aa):536

Product Details

Purity: > 90 % by SDS-PAGE

Target Details

Target: FLT1

Alternative Name: VEGFR-1/Flt-1 ([FLT1 Products](#))

Background: Recombinant human soluble Vascular Endothelial Growth Factor Receptor-1 domain D1-5 (sVEGFR-1(D5)) is a 70 kDa protein. The baculovirus generated, recombinant human sVEGFR-1 is produced as a non-chimeric protein in a monomeric form. The soluble receptor protein contains only the first 5 extracellular domains, which contain all the information necessary for high affinity ligand binding. The receptor monomers have a mass of approximately 70 kDa. Endothelial cells express three different vascular endothelial growth factor (VEGF) receptors, belonging to the family of receptor tyrosine kinases (RTKs). They are named VEGFR-1 (Flt-1), VEGFR-2 (KDR/Flk-1), VEGFR-3 (Flt-4). Their expression is almost exclusively restricted to endothelial cells, but VEGFR-1 can also be found on monocytes, dendritic cells and on trophoblast cells. The flt-1 gene was first described in 1990. The receptor contains seven immunoglobulin-like extracellular domains, a single transmembrane region and an intracellular split tyrosine kinase domain. Compared to VEGFR-2 the Flt-1 receptor has a higher affinity for VEGF but a weaker signaling activity. VEGFR-1 thus leads not to proliferation of endothelial cells, but mediates signals for differentiation. Interestingly a naturally occurring soluble variant of VEGFR-1 (sVEGFR-1) was found in HUVEC supernatants in 1996, which is generated by alternative splicing of the flt-1 mRNA. The biological functions of sVEGFR-1 still are not clear, but it seems to be an endogenous regulator of angiogenesis, binding VEGF with the same affinity as the full-length receptor.

Synonyms: soluble vascular endothelial growth factor receptor-1, soluble FLT1, soluble VEGFR-1

Molecular Weight: 70.0 kDa

Gene ID: 2321

NCBI Accession: [NM_001159920](#), [NP_001153392](#)

UniProt: [P17948](#)

Pathways: [RTK Signaling](#), [Signaling Events mediated by VEGFR1 and VEGFR2](#), [VEGFR1 Specific Signals](#)

Application Details

Application Notes: The activity of sVEGFR-1(D5) was determined by its ability to inhibit the VEGF-A-induced proliferation of HUVECs.

Comment: Soluble Receptors

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: The lyophilized human sVEGFR-1(D5) is soluble in water and most aqueous buffers. The lyophilized powder should be reconstituted in water to a concentration not lower than 100 μ g/mL.

Buffer: PBS

Storage: -20 °C, -80 °C

Storage Comment: Lyophilized samples are stable for greater than six months at -20°C to -70°C. Reconstituted sVEGFR-1(D5) should be stored in working aliquots at -70°C.

Expiry Date: 6 months