

Datasheet for ABIN1589899
anti-EGFL7 antibody (C-Term)



[Go to Product page](#)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µg |
| Target: | EGFL7 |
| Binding Specificity: | C-Term |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This EGFL7 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

Product Details

| | |
|------------------|--|
| Purpose: | VE-Statin/EGFL7 antibody |
| Immunogen: | Recombinant human EGFL7 C-Terminus |
| Isotype: | IgG |
| Specificity: | Recombinant human EGFL-7 C-Terminus |
| Characteristics: | Chromosomal location: 9q34.3 Produced from sera of rabbits immunised with highly pure recombinant human VE-Statin/EGFL-7. |
| Purification: | Anti-human VE-Statin/EGFL-7 was purified by Protein-A chromatography. |

Target Details

| | |
|-------------------|---|
| Target: | EGFL7 |
| Alternative Name: | VE-Statin/EGFL7 (EGFL7 Products) |
| Background: | <p>EGFL7, NEU1, ZNEU1, VE-STATIN, RP11-251M1.2, NOTCH4-like protein, Vascular endothelial statin,EGFL7 (VE-Statin) is an ~ 30 kDa secreted protein that contains an Emilin-like (EMI) domain (a multimerization motif), and two epidermal growth factor (EGF) domains, one of which binds calcium. Based on these domains, it has been hypothesized that EGFL7 may self-assemble like extracellular matrix (ECM) proteins and, thus, could incorporate into ECM. EGFL7 has been reported to stimulate cell adhesion as well as motility in a manner similar to ECM proteins. EGFL7 has been shown to be primarily expressed by developing ECs but also by primordial germ cells and some central nervous system neurons. Interestingly, EGFL7 expression markedly decreases in ECs in postnatal life, but can be strongly up-regulated after various tissue injuries that lead to increased angiogenic responses.</p> |
| Gene ID: | 51164 |
| NCBI Accession: | NM_016215 , NP_057299 |
| UniProt: | Q9UHF1 |

Application Details

| | |
|--------------------|-----------------------------|
| Application Notes: | Western Blot: Use 2-5 µg/mL |
| Restrictions: | For Research Use only |

Handling

| | |
|------------------|---|
| Format: | Lyophilized |
| Reconstitution: | Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/mL. |
| Buffer: | PBS |
| Handling Advice: | Centrifuge vial prior to opening. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | The lyophilized antibody is stable for at least 2 years at -20°C. After sterile reconstitution the antibody is stable at 2-8°C for up to 6 months. Frozen aliquots are stable for at least 6 months when stored at -20°C. Addition of a carrier protein or 50% glycerol is recommended for frozen aliquots. |

Handling

Expiry Date: 24 months