

[Go to Product page](#)

Datasheet for ABIN1589965

TrueBlot® Anti-Mouse Ig IP Beads

Overview

Quantity:	2.5 mL
Reactivity:	Mouse
Host:	Goat
Clonality:	Polyclonal
Conjugate:	Agarose Beads
Application:	Immunoprecipitation (IP), Western Blotting (WB)

Product Details

Brand:	TrueBlot®
Characteristics:	<p>TrueBlot® anti-Mouse Ig IP Beads are a suspension of activated agarose beads coupled with goat anti-mouse IgG. It is suitable for precipitation of mouse IgGs used as the primary antibodies in immunoprecipitation assays. The beads are in suspension and will settle upon storage. Prior to use, mix the vial gently (do not vortex) to ensure delivery of proper bead volume.</p> <p>Conjugation Name: Agarose beads for TrueBlot®</p>
Components:	TrueBlot® Anti-Mouse Ig IP Beads

Application Details

Application Notes:	<p>Immunoprecipitation Dilution: TrueBlot® anti-Mouse Ig IP Beads (binds 0.4 mg Ig/mL beads) have been reported for use in IP</p> <p>Western Blot Dilution: Use with Mouse TrueBlot® (ABIN1589977)</p>
Comment:	Upon initial use of this product, we recommend that the vial be inverted several times to get the

Application Details

beads into suspension. We recommend to use a large bore pipet to pipet up the liquid for use. For storage of the opened vial, we recommend that the vial cap be sealed with parafilm to help prevent evaporation of the buffer.

Assay Procedure:	<p>Preparation of Immunoprecipitated Sample for SDS-PAGE:</p> <p>1. Preclear cell lysate: Add 50 µL of Anti-Mouse IgG Beads and 500µl of cell lysate sample to a microcentrifuge tube and incubate on ice for</p> <p>3. minutes. Spin at 10,000xg for 3 minutes and transfer the supernatant to a new microcentrifuge tube.</p> <p>2. Immunoprecipitation: Add 5 µg of primary antibody to the microcentrifuge tube containing the precleared lysate. Incubate on ice for 1 hour. Add 50 µL of Anti-Mouse IgG Beads. Incubate for 1 hour on a rocking platform. Spin the microcentrifuge tube at 10000xg for 1 minute. Remove supernatant completely and wash the (pelleted) beads 3 times with 500 µL of Lysis Buffer (50mM Tris HCl pH 8.0, 150mM NaCl,1 % NP-40).</p> <p>3. Prepare sample for SDS-PAGE: After the last wash, aspirate supernatant, and add 100 µL Laemmli Buffer (with 50 mM DTT or 1 %β-mercaptoethanol, final) to bead pellet. Vortex and heat to 90-100 °C for</p> <p>1. minutes. Spin at 10000xg for 3 minutes, collect supernatant, and load onto the gel. Avoid loading Anti-Mouse Ig Beads.</p> <p>Note: The supernatant can be stored at -20 °C for future use. After thawing, add fresh dithiothreitol and heat as above. Centrifuge the sample at 10000xg for 1 minute in a microcentrifuge tube to pellet any Anti-Mouse Ig Beads and immediately transfer an aliquot of the supernatant to gel wells.</p>
------------------	---

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
Buffer:	<p>Buffer: 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2</p> <p>0.01 % (w/v) Sodium Azide</p> <p>Stabilizer: None</p>
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Do not freeze.

Handling

Sensitive to light.
Prior to use, mix the vial gently (do not vortex) to ensure delivery of proper bead volume.

Storage: 4 °C

Storage Comment: Store vial at 4 °C prior to opening.

Expiry Date: 6 months