antibodies -online.com





Datasheet for ABIN1589966

TrueBlot® Anti-Goat Ig IP Beads



| () | ۱ ۱ | \cap | r | /1 | \cap | ۱ ۸ | 1 |
|-----|-----|--------|----|----|--------|-----|---|
| 0 | 'V | ㄷ | I١ | νı | \Box | V | ۷ |

| Quantity: | 2.5 mL |
|--------------|---|
| Reactivity: | Goat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | Agarose Beads |
| Application: | Immunoprecipitation (IP), Western Blotting (WB) |

Product Details

| Brand: | TrueBlot® |
|------------------|--|
| Characteristics: | TrueBlot® anti-Goat Ig IP Beads are a suspension of activated agarose beads coupled with rabbit anti-goat IgG. It is suitable for precipitation of goat IgGs used as the primary antibodies in |
| | immunoprecipitation assays. The beads are in suspension and will settle upon storage. Prior to |
| | use, mix the vial gently (do not vortex) to ensure delivery of proper bead volume. Conjugation Name: Agarose beads for TrueBlot® |
| Components: | TrueBlot® Anti-Goat Ig IP Beads |

Application Details

| Application Notes: | Immunoprecipitation Dilution: TrueBlot® anti-Goat Ig IP Beads (binds 1 mg Ig/mL beads) have | | |
|--------------------|---|--|--|
| | been reported for use in IP | | |
| | Western Blot Dilution: Use with Goat TrueBlot® (ABIN1589970) | | |
| Comment: | Upon initial use of this product, we recommend that the vial be inverted several times to get the | | |
| | beads into suspension. We recommend to use a large bore pipet to pipet up the liquid for use. | | |

Application Details

| | For storage of the opened vial, we recommend that the vial cap be sealed with parafilm to help |
|--------------------|--|
| | prevent evaporation of the buffer. |
| Assay Procedure: | Preparation of Immunoprecipitated Sample for SDS-PAGE: |
| | 1. Preclear cell lysate: Add 50 μL of anti-goat IgG beads and 500 μL of cell lysate sample to an |
| | eppendorf tube and incubate on ice for |
| | 3. minutes. Spin at 10,000xg for 3 minutes and transfer the supernatant to a new eppendorf |
| | tube. |
| | 2. Immunoprecipitation: Add 5 μg of primary antibody to the eppendorf tube containing the |
| | precleared lysate. Incubate on ice for 1 hour. Add 50 µL of Anti-Goat IgG Beads. Incubate for 1 |
| | hour on a rocking platform. Spin the eppendorf tube at 10000xg for 1 minute. Remove |
| | supernatant completely and wash the (pelleted) beads 3 times with 500 μL of Lysis Buffer. |
| | 3. Prepare sample for SDS-PAGE: After the last wash, aspirate supernatant, and add 100 μ L |
| | Laemmli Buffer (with 50 mM DTT or 2 % ß-mercaptoethanol, final) to bead pellet. Vortex and |
| | heat to 90-100 °C for |
| | 1. minutes. Spin at 10000xg for 3 minutes, collect supernatant, and load onto the gel. Avoid |
| | loading anti-goat Ig beads. |
| | Note: The supernatant can be stored at -20 °C for future use. After thawing, add fresh Reducin |
| | Agent (dithiothreitol) and heat as above. Centrifuge the sample at 10000xg for 1 minute in a |
| | microcentrifuge to pellet any anti-goat Ig beads and immediately transfer an aliquot of the |
| | supernatant to gel wells. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Buffer: | Buffer: 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| | 0.01 % (w/v) Sodium Azide |
| | Stabilizer: None |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which |
| | should be handled by trained staff only. |
| Handling Advice: | Do not freeze. |
| | Sensitive to light. |
| | Prior to use, mix the vial gently (do not vortex) to ensure delivery of proper bead volume. |

Handling

| Storage: | 4 °C |
|------------------|--------------------------------------|
| Storage Comment: | Store vial at 4 °C prior to opening. |
| Expiry Date: | 6 months |