# antibodies -online.com





# anti-TRAF3IP2 antibody (Internal Region)





Go to Product page

_					
U	V	er	VI	е	W

Quantity:	100 μg
Target:	TRAF3IP2
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This TRAF3IP2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

#### **Product Details**

Purpose:	TRAF3IP2	
Sequence:	PDTGHDSDKS DQ	
Isotype:	IgG	
Specificity:	This antibody is expected to recognise reported isoforms 2 and 3 (NP_679211.1, NP_001157753.1).	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.	
Grade:	Verified	

## **Target Details**

rarget Details		
Target:	TRAF3IP2	
Alternative Name:	TRAF3IP2 (TRAF3IP2 Products)	
Background:	TRAF3IP2, TRAF3 interacting protein 2 , ACT1, C6orf4, C6orf5, C6orf6, CIKS, DKFZP586G0522	
	MGC3581, NFkB-activating protein ACT1, OTTHUMP0000040422, connection to IKK and SAPK/JNK	
	SAPR/JINK	
Gene ID:	10758	
NCBI Accession:	NP_679211, NP_001157753	
Pathways:	Production of Molecular Mediator of Immune Response	
Application Details		
Application Notes:	Western Blot: Approx 65 kDa band observed in lysates of cell line MCF7 (calculated MW of	
	63.6 kDa according to NP_679211.1). Recommended concentration: 1-3 $\mu$ g/mL.	
	Peptide ELISA: antibody detection limit dilution 1:2000.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	0.5 mg/mL	
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum	
	albumin.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	
Handling Advice:	Minimize freezing and thawing.	
Storage:	-20 °C	
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated	
	at 4°C for a few weeks and still remain viable.	

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

### **Western Blotting**

**Image 1.** ABIN1590027 (2μg/ml) staining of MCF7 lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.