

Datasheet for ABIN1607709

anti-Dextranase antibody (Biotin)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µg
Target:	Dextranase
Reactivity:	Penicillium sp.
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Dextranase antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunoprecipitation (IP)

Product Details

Purpose:	Dextranase (Penicillium Species) Antibody Biotin Conjugated
Immunogen:	Immunogen: Dextranase [Penicillium species] DEX Immunogen Type: Native Protein
Isotype:	IgG
Cross-Reactivity (Details):	Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Rabbit Serum as well as purified and partially purified Dextranase [Penicillium species].
Characteristics:	Synonyms: rabbit anti-Dextranase Antibody biotin Conjugation, biotin Conjugated rabbit anti-Dextranase Antibody, dextranase, Alpha-1,6-glucan-6-glucanohydrolase, DEX
Purification:	Anti-Dextranase is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above.

Target Details

Target:	Dextranase
Abstract:	Dextranase Products
Background:	<p>Background: Anti-Dextranase antibody is a transferase enzyme which catalyzes the synthesis of dextran by engaging in the endohydrolysis of 1, 6 alpha-D-glucosidic linkages in dextran.</p> <p>Belonging to the glycosyl hydrolase 49 family, Anti-Dextranase antibody is ideal for researchers interested in Metabolism and Signal Transduction research.</p>
UniProt:	P48845

Application Details

Application Notes:	<p>Application Note: Anti-Dextranase Biotin Antibody has been tested by ELISA and western blot and is assayed against 1.0 µg of Dextranase in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:3,000 to 1:14,000 of the reconstitution concentration is suggested for this product.</p> <p>Western Blot Dilution: 1:500 - 1:5,000</p> <p>Immunoprecipitation Dilution: 1:100</p> <p>ELISA Dilution: 1:2,000 - 1:10,000</p> <p>Other: User Optimized</p>
Restrictions:	For Research Use only


Handling

Format:	Lyophilized
Reconstitution:	<p>Reconstitution Volume: 100 µL</p> <p>Reconstitution Buffer: Restore with deionized water (or equivalent)</p>
Concentration:	10 mg/mL
Buffer:	<p>Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2</p> <p>Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free</p> <p>Preservative: 0.01 % (w/v) Sodium Azide</p>
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling


Storage:	4 °C,-20 °C
Storage Comment:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiry Date:	12 months

Images



ELISA

Image 1. ELISA Results of Rabbit Anti-Dextranase Antibody Biotin Conjugated tested against purified Dextranase protein. Each well was coated in duplicate with 1.0 µg of Dextranase. The starting dilution of antibody was 5 µg/mL and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using HRP Conjugate Stabilizer (p/n MB-076), Streptavidin-HRP conjugated (p/n S000-03), and TMB substrate (p/n TMBE-1000).



Western Blotting

Image 2. Western Blot of Anti-Dextranase Antibody Biotin Conjugated. Lane 1: Molecular Weight Marker. Lane 2: Dextranase. Primary Antibody: Rabbit Anti-Dextranase Biotin Conjugate at 1:1000 overnight at 2-8 °C. Secondary Antibody: Streptavidin-HRP (p/n S000-03) at 1:40,000 for 30 min at RT. Block: BlockOut Buffer (p/n MB-073). Expected MW: ~66 kDa.