antibodies -online.com





Pleckstrin Protein (PLEK) (AA 1-350) (His tag)



Overview

1 mg
Pleckstrin (PLEK)
AA 1-350
Rat
Yeast
Recombinant
This Pleckstrin protein is labelled with His tag.
ELISA

Product Details	
Sequence:	MEPKRIREGY LVKKGSVFNT WKPMWVVLLE DGIEFYKKKS DNNPKGMIPL KGSTLTSPCQ DFGKRMFVLK ITTTKQQDHF FQAAYLEERD AWVRDIKKAI KCIEGGQKFA RKSTRRSIRL PETIDLGALY LSMKDPEKGI KELNLEKDKK VFNHCFTGSG VIDWLVSNKL VRNRQEGLMI SASLLSEGYL QPASDLSKNA ADGIAENPFL DNPDAFYYFP DSGFFCEENS SDDDIILKEE FRGIIIKQGC LLKQGHRRKN WKVRKFILRE DPAYLHYYDP AGGEDPLGAI HLRGCVVTSV ESNPDGKKSD EENLFEIITA DEVHYYMQAA TAKERTEWIK AIQVASRTGK
Specificity:	Rattus norvegicus (Rat)
Characteristics:	Please inquire if you are interested in this recombinant protein expressed in E. coli, mammalien cells or by baculovirus infection. Be aware about differences in price and lead time.
Purity:	> 90 %

Target Details

Target:	Pleckstrin (PLEK)
Abstract:	PLEK Products
Background:	Recommended name: Pleckstrin
UniProt:	Q4KM33
Pathways:	Inositol Metabolic Process, Regulation of G-Protein Coupled Receptor Protein Signaling, Regulation of Cell Size, Regulation of Carbohydrate Metabolic Process

Application Details

Comment:

The yeast protein expression system is the most economical and efficient eukaryotic system for secretion and intracellular expression. A protein expressed by the mammalian cell system is of very high-quality and close to the natural protein. But the low expression level, the high cost of medium and the culture conditions restrict the promotion of mammalian cell expression systems. The yeast protein expression system serve as a eukaryotic system integrate the advantages of the mammalian cell expression system. A protein expressed by yeast system could be modificated such as glycosylation, acylation, phosphorylation and so on to ensure the native protein conformation. It can be used to produce protein material with high added value that is very close to the natural protein. Our proteins produced by yeast expression system has been used as raw materials for downstream preparation of monoclonal antibodies.

Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Concentration:	0.2-2 mg/mL
Buffer:	Tris-based buffer, 50 % glycerol
Handling Advice:	Repeated freezing and thawing is not recommended. Store working aliquots at 4 °C for up to one week
Storage:	-20 °C
Storage Comment:	Store at -20 °C, for extended storage, conserve at -20 °C or -80 °C.