# antibodies -online.com







## NACA Protein (AA 1-215) (His tag)



$\sim$			
	N/P	r\/	i⊢₩

1 mg	
NACA (NACa1)	
AA 1-215	
Cow	
Yeast	
Recombinant	
This NACA protein is labelled with His tag.	
ELISA	
MPGEATDTVP ATEQELPQPQ AETGSGTESD SDESVPELEE QDSTQATTQQ AQLAAAAEID	
EEPVSKAKQS RSEKKARKAM SKLGLRQVTG VTRVTIRKSK NILFVITKPD VYKSPASDTY	
IVFGEAKIED LSQQAQLAAA EKFKVQGEAV SNIQENTQTP TVQEESEEEE VDETGVEVKD	
IELVMSQANV SRAKAVRALK NNSNDIVNAI MELTM	
Bos taurus (Bovine)	
Please inquire if you are interested in this recombinant protein expressed in E. coli, mammalien	
cells or by baculovirus infection. Be aware about differences in price and lead time.	
> 90 %	
NACA (NACa1)	

#### **Target Details**

Alternative Name:	Nascent polypeptide-associated complex subunit alpha (NACA) (NACa1 Products)	
Background:	Recommended name: Nascent polypeptide-associated complex subunit alpha.  Short name= NAC-alpha.  Alternative name(s): Alpha-NAC	
UniProt:	Q5E9A1	
Pathways:	Regulation of Muscle Cell Differentiation, Skeletal Muscle Fiber Development	

#### **Application Details**

#### Comment:

The yeast protein expression system is the most economical and efficient eukaryotic system for secretion and intracellular expression. A protein expressed by the mammalian cell system is of very high-quality and close to the natural protein. But the low expression level, the high cost of medium and the culture conditions restrict the promotion of mammalian cell expression systems. The yeast protein expression system serve as a eukaryotic system integrate the advantages of the mammalian cell expression system. A protein expressed by yeast system could be modificated such as glycosylation, acylation, phosphorylation and so on to ensure the native protein conformation. It can be used to produce protein material with high added value that is very close to the natural protein. Our proteins produced by yeast expression system has been used as raw materials for downstream preparation of monoclonal antibodies.

Restrictions:

For Research Use only

### Handling

Format:	Lyophilized	
Concentration:	0.2-2 mg/mL	
Buffer:	Tris-based buffer, 50 % glycerol	
Handling Advice:	Repeated freezing and thawing is not recommended. Store working aliquots at 4 °C for up to one week	
Storage:	-20 °C	
Storage Comment:	Store at -20 °C, for extended storage, conserve at -20 °C or -80 °C.	