antibodies .- online.com





Fatty Acid/Retinol Binding Protein Family Member (Far-1) (FAR-1) (AA 17-178) protein (His tag)



Go to Product page

_	
	rview
	VICVV

1 mg
Fatty Acid/Retinol Binding Protein Family Member (Far-1) (FAR-1)
AA 17-178
Wuchereria bancrofti
Yeast
Recombinant
His tag
ELISA
NVIP FSMSTIPEEY KEFIPEEVRN FYKDLTVEDK EILRELASKH ATFANEDAAL EALKAKSDNL
YKNAVELRNF VKAKIDSLKP DAKTFVDEII AKARSLRSDD GHKLDTEKIK QAARDIIAKY
QALSEETKEE LKVTFPAIAK IIGNEKLKRN ASTFLQKN
Wuchereria bancrofti
Please inquire if you are interested in this recombinant protein expressed in E. coli, mammalien
cells or by baculovirus infection. Be aware about differences in price and lead time.
> 90 %
Fatty Acid/Retinol Binding Protein Family Member (Far-1) (FAR-1)
• • • • • • • • • • • • • • • • • • •

Target Details

Background:	Recommended name: Fatty-acid and retinol-binding protein 1.	
	Alternative name(s): Wb-FAR-1	
UniProt:	Q8WT54	

Application Details

Comment:

The yeast protein expression system is the most economical and efficient eukaryotic system for secretion and intracellular expression. A protein expressed by the mammalian cell system is of very high-quality and close to the natural protein. But the low expression level, the high cost of medium and the culture conditions restrict the promotion of mammalian cell expression systems. The yeast protein expression system serve as a eukaryotic system integrate the advantages of the mammalian cell expression system. A protein expressed by yeast system could be modificated such as glycosylation, acylation, phosphorylation and so on to ensure the native protein conformation. It can be used to produce protein material with high added value that is very close to the natural protein. Our proteins produced by yeast expression system has been used as raw materials for downstream preparation of monoclonal antibodies.

Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Concentration:	0.2-2 mg/mL
Buffer:	Tris-based buffer, 50 % glycerol
Handling Advice:	Repeated freezing and thawing is not recommended. Store working aliquots at 4 °C for up to one week
Storage:	-20 °C
Storage Comment:	Store at -20 °C, for extended storage, conserve at -20 °C or -80 °C.