
antibodies .- online.com





PSMG1 Protein (AA 1-257) (His tag)

> 90 %



()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Purity:

OVEIVIEW	
Quantity:	1 mg
Target:	PSMG1
Protein Characteristics:	AA 1-257
Origin:	Nematostella vectensis
Source:	Yeast
Protein Type:	Recombinant
Purification tag / Conjugate:	This PSMG1 protein is labelled with His tag.
Application:	ELISA
Product Details	
Sequence:	MVDAFGLWEL NFPCSRVDDD DDEEEQITKE QEEQLCPQFV FSPQFSKDVG KTEPIECSVL VLGVGEVAST FLEVHFLCGN NSSVAAVISD KKTEKNFEYL VKNLRNKRSS LLHQLTRNND VMVCQISSHV EQDKGFYWTQ KIFSHIRPAK VVILTSSPAS EYNSDDPSEI KTDFVKMMKT NAWSNSCAGP DITDIQTPNT VKGLAASVLT HCQIYKLPAA LFMCYTESMH VDAQAVEAFR CKAPYKQSSM GLESGKL
Specificity:	Nematostella vectensis (Starlet sea anemone)
Characteristics:	Please inquire if you are interested in this recombinant protein expressed in E. coli, mammalien cells or by baculovirus infection. Be aware about differences in price and lead time.
D ::	00.0

Target Details

Target:	PSMG1
Alternative Name:	Proteasome assembly chaperone 1 (psmg1) (PSMG1 Products)
Background:	Recommended name: Proteasome assembly chaperone 1
UniProt:	A7RV13

Application Details

Comment:

The yeast protein expression system is the most economical and efficient eukaryotic system for secretion and intracellular expression. A protein expressed by the mammalian cell system is of very high-quality and close to the natural protein. But the low expression level, the high cost of medium and the culture conditions restrict the promotion of mammalian cell expression systems. The yeast protein expression system serve as a eukaryotic system integrate the advantages of the mammalian cell expression system. A protein expressed by yeast system could be modificated such as glycosylation, acylation, phosphorylation and so on to ensure the native protein conformation. It can be used to produce protein material with high added value that is very close to the natural protein. Our proteins produced by yeast expression system has been used as raw materials for downstream preparation of monoclonal antibodies.

Restrictions:

For Research Use only

Handling

Format:	Lyophilized	
Concentration:	0.2-2 mg/mL	
Buffer:	Tris-based buffer, 50 % glycerol	
Handling Advice:	Repeated freezing and thawing is not recommended. Store working aliquots at 4 °C for up to one week	
Storage:	-20 °C	
Storage Comment:	Store at -20 °C, for extended storage, conserve at -20 °C or -80 °C.	