

Datasheet for ABIN1655621 NAGK Protein (AA 1-302) (His tag)



Overview

Overview	
Quantity:	1 mg
Target:	NAGK
Protein Characteristics:	AA 1-302
Origin:	Vibrio splendidus
Source:	Yeast
Protein Type:	Recombinant
Purification tag / Conjugate:	This NAGK protein is labelled with His tag.
Application:	ELISA
Product Details	
Sequence:	MYYGFDVGGT KIEFGAFNEK LERVATERVP TPTEDYQLLL DTIAGLVKKY DNEFSCEGKI
	GLGLPGMEDA DDGTMLVVNV PASTGKPLRK DLEALIGRSV KIENDANCFA LSEAWDDELK
	DEPSVAGLIL GTGFGGGLVY EGKVFSGRNH VAGELGHMRL PLDAWFHLGD NAPLLGCGCG
	KKGCLDSYLS GRGFELIYEH YFGEKKKAIE IIQAYNEGES KAAEHVDRFM ELLAICFANL
	FTGLDPHVVA LGGGLSNFEL IYEEMPKRIP KYLLSVAKCP KIIKAKHGDS GGVRGAAFLN IK
Specificity:	Vibrio splendidus (strain LGP32) (Vibrio splendidus (strain Mel32))
Characteristics:	Please inquire if you are interested in this recombinant protein expressed in E. coli, mammalien
	cells or by baculovirus infection. Be aware about differences in price and lead time.
Purity:	> 90 %

Target Details

Target:	NAGK
Alternative Name:	N-acetyl-D-glucosamine kinase (nagK) (NAGK Products)
Background:	Recommended name: N-acetyl-D-glucosamine kinase. EC= 2.7.1.59. Alternative name(s): GlcNAc kinase
UniProt:	B7VNU4

Application Details

Comment:

The yeast protein expression system is the most economical and efficient eukaryotic system for secretion and intracellular expression. A protein expressed by the mammalian cell system is of very high-quality and close to the natural protein. But the low expression level, the high cost of medium and the culture conditions restrict the promotion of mammalian cell expression systems. The yeast protein expression system serve as a eukaryotic system integrate the advantages of the mammalian cell expression system. A protein expressed by yeast system could be modificated such as glycosylation, acylation, phosphorylation and so on to ensure the native protein conformation. It can be used to produce protein material with high added value that is very close to the natural protein. Our proteins produced by yeast expression system has been used as raw materials for downstream preparation of monoclonal antibodies.

Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Concentration:	0.2-2 mg/mL
Buffer:	Tris-based buffer, 50 % glycerol
Handling Advice:	Repeated freezing and thawing is not recommended. Store working aliquots at 4 °C for up to one week
Storage:	-20 °C
Storage Comment:	Store at -20 °C, for extended storage, conserve at -20 °C or -80 °C.