





Ndd Nucleoid Disruption Protein (NDD) (AA 1-151) protein (His tag)



Go to Product page

\sim						
	1//	\Box	r۱	/1	0	Λ

Quantity:	1 mg
Target:	Ndd Nucleoid Disruption Protein (NDD)
Protein Characteristics:	AA 1-151
Origin:	Enterobacteria phage T6
Source:	Yeast
Protein Type:	Recombinant
Purification tag / Conjugate:	His tag
Application:	ELISA
Product Details	
Sequence:	MKYMTVTDLN NAGATVIGTI KGGEWFLGTP HKDILSKPGF YFLVSKLGGP FSSPCVSARF YVGNQRSKQG FSAVLSHIRQ RRSQLARTIA NNNVPYTVFY LPASKMKPLT TGFGKGQLAL AFTRNHHSEY QTLEEMNRML ADNFKFVLQA Y
Specificity:	Enterobacteria phage T6 (Bacteriophage T6)
Characteristics:	Please inquire if you are interested in this recombinant protein expressed in E. coli, mammalien cells or by baculovirus infection. Be aware about differences in price and lead time.
Purity:	> 90 %
Target Details	
Target:	Ndd Nucleoid Disruption Protein (NDD)

Target Details

Target Type:	Phage Protein	
Background:	Recommended name: Nuclear disruption protein	
UniProt:	P42265	

Application Details

Comment:

The yeast protein expression system is the most economical and efficient eukaryotic system for secretion and intracellular expression. A protein expressed by the mammalian cell system is of very high-quality and close to the natural protein. But the low expression level, the high cost of medium and the culture conditions restrict the promotion of mammalian cell expression systems. The yeast protein expression system serve as a eukaryotic system integrate the advantages of the mammalian cell expression system. A protein expressed by yeast system could be modificated such as glycosylation, acylation, phosphorylation and so on to ensure the native protein conformation. It can be used to produce protein material with high added value that is very close to the natural protein. Our proteins produced by yeast expression system has been used as raw materials for downstream preparation of monoclonal antibodies.

Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Concentration:	0.2-2 mg/mL
Buffer:	Tris-based buffer, 50 % glycerol
Handling Advice:	Repeated freezing and thawing is not recommended. Store working aliquots at 4 °C for up to one week
Storage:	-20 °C
Storage Comment:	Store at -20 °C, for extended storage, conserve at -20 °C or -80 °C.