# antibodies -online.com





## Hemoglobin, beta Adult Minor Chain (HBB-B2) (AA 2-146) protein (His tag)



Go to Product page

0	1 /	0	K	/ 1	$\sim$	1 /	١.
	1//	$\vdash$	ı ۱	71	$\vdash$	W	W

Alternative Name:

1 mg	
Hemoglobin, beta Adult Minor Chain (HBB-B2)	
AA 2-146	
Xenopus borealis	
Yeast	
Recombinant	
His tag	
ELISA	
GLTAHEKQL ITGSWGKINA KAIGKEALGR LLNTFPWTQR YFSSFGNLGS AEAIFHNEAV	
AAHGEKVVTS VGEAIKHMDD IKGYYAELSK YHSETLHVDP NNFKRFGGCL SITLGHHFGE	
EYTPELHAAY EHLFDAIADA LGKGYH	
Xenopus borealis (Kenyan clawed frog)	
Please inquire if you are interested in this recombinant protein expressed in E. coli, mammalien	
cells or by baculovirus infection. Be aware about differences in price and lead time.	
> 90 %	
Hemoglobin, beta Adult Minor Chain (HBB-B2)	

Hemoglobin subunit beta-2 (hbb2) (HBB-B2 Products)

#### **Target Details**

Background:	Recommended name: Hemoglobin subunit beta-2.		
	Alternative name(s): Beta-2-globin Hemoglobin beta-2 chain Hemoglobin beta-minor chain		
UniProt:	P07433		

### **Application Details**

#### Comment:

The yeast protein expression system is the most economical and efficient eukaryotic system for secretion and intracellular expression. A protein expressed by the mammalian cell system is of very high-quality and close to the natural protein. But the low expression level, the high cost of medium and the culture conditions restrict the promotion of mammalian cell expression systems. The yeast protein expression system serve as a eukaryotic system integrate the advantages of the mammalian cell expression system. A protein expressed by yeast system could be modificated such as glycosylation, acylation, phosphorylation and so on to ensure the native protein conformation. It can be used to produce protein material with high added value that is very close to the natural protein. Our proteins produced by yeast expression system has been used as raw materials for downstream preparation of monoclonal antibodies.

Restrictions:

For Research Use only

#### Handling

Format:	Lyophilized	
Concentration:	0.2-2 mg/mL	
Buffer:	Tris-based buffer, 50 % glycerol	
Handling Advice:	Repeated freezing and thawing is not recommended. Store working aliquots at 4 °C for up to one week	
Storage:	-20 °C	
Storage Comment:	Store at -20 °C, for extended storage, conserve at -20 °C or -80 °C.	