



[Go to Product page](#)

Datasheet for ABIN1671935

Carkd Protein (AA 1-303) (His tag)

Overview

Quantity:	1 mg
Target:	Carkd
Protein Characteristics:	AA 1-303
Origin:	Tick
Source:	Yeast
Protein Type:	Recombinant
Purification tag / Conjugate:	This Carkd protein is labelled with His tag.
Application:	ELISA

Product Details

Sequence:	MVPHVPEGCS VQQQLVCSVI PPLNSERRKG QAGRVGIVGG SAEYTGAPYF AAMAALRTGA DLVHVFCHPS AATAIKAYSP ELIVHPTLDA AVTCLPRLHA VVVGPGGLGRD VEASWMPTLF NRIREQGLPV VVDADGLFYV TQNPDLVRGY SRAILTPNAV ELDRLYRAVL GSPPRENAV ELARALGHVT VLAKGSEDII SDGHRLLRCT EQGSPRRCGG QGDLVSGSLA LFAFWSHSAH DTPGEASKRQ NSEYGPAMIA ALGAAMLVRR CGRLAFQKMA RSTLSSDMLA EVRTAFSMLF PVD
Specificity:	Ixodes scapularis (Black-legged tick) (Deer tick)
Characteristics:	Please inquire if you are interested in this recombinant protein expressed in E. coli, mammalian cells or by baculovirus infection. Be aware about differences in price and lead time.
Purity:	> 90 %

Target Details

Target:	Carkd
Alternative Name:	ATP-dependent (S)-NAD (P)H-hydrate dehydratase (Carkd Products)
Background:	Recommended name: ATP-dependent (S)-NAD(P)H-hydrate dehydratase. EC= 4.2.1.93. Alternative name(s): ATP-dependent NAD(P)HX dehydratase
UniProt:	B7PBI5

Application Details

Comment:	The yeast protein expression system is the most economical and efficient eukaryotic system for secretion and intracellular expression. A protein expressed by the mammalian cell system is of very high-quality and close to the natural protein. But the low expression level, the high cost of medium and the culture conditions restrict the promotion of mammalian cell expression systems. The yeast protein expression system serve as a eukaryotic system integrate the advantages of the mammalian cell expression system. A protein expressed by yeast system could be modiflicated such as glycosylation, acylation, phosphorylation and so on to ensure the native protein conformation. It can be used to produce protein material with high added value that is very close to the natural protein. Our proteins produced by yeast expression system has been used as raw materials for downstream preparation of monoclonal antibodies.
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Concentration:	0.2-2 mg/mL
Buffer:	Tris-based buffer, 50 % glycerol
Handling Advice:	Repeated freezing and thawing is not recommended. Store working aliquots at 4 °C for up to one week
Storage:	-20 °C
Storage Comment:	Store at -20 °C, for extended storage, conserve at -20 °C or -80 °C.