antibodies .- online.com





Dre2p (DRE2) (AA 1-341) protein (His tag)



Overview

Quantity:	1 mg
Target:	Dre2p (DRE2)
Protein Characteristics:	AA 1-341
Origin:	Pichia pastoris
Source:	Yeast
Protein Type:	Recombinant
Purification tag / Conjugate:	His tag
Application:	ELISA

Product Details

IVTEPQAVEA AKDSLAQKLG TDKSNISQHI IDRVATGQVS LRPSYYNHIQ
KLPVKCYEII FEALQSNGVF EGTIPPESAT DGILQGFLVE SDTKWVKPST
AGQNKTKSSA FKKEMPFFKK LSSPPTLTDS SEADEDEESQ LNEKLKGSKL
DEDELLRD DDGALKGPVV VPVKCALPNG KRRKKACKDC TCGLKELEEN
SILSKLATSA NAEAEKIEER LRRKQASKEG EEVKFTEQEV TEIDFTIQGK
DAFRCDGCPY LGLPPFKPGQ AISIEGLGAD I
(strain GS115 / ATCC 20864) (Yeast)
if you are interested in this recombinant protein expressed in E. coli, mammalien
ulovirus infection. Be aware about differences in price and lead time.
,

Target Details

Target:	Dre2p (DRE2)
Alternative Name:	Fe-S cluster assembly protein DRE2 (DRE2) (DRE2 Products)
Background:	Recommended name: Fe-S cluster assembly protein DRE2. Alternative name(s): Anamorsin homolog
UniProt:	C4QY17

Application Details

Comment:

The yeast protein expression system is the most economical and efficient eukaryotic system for secretion and intracellular expression. A protein expressed by the mammalian cell system is of very high-quality and close to the natural protein. But the low expression level, the high cost of medium and the culture conditions restrict the promotion of mammalian cell expression systems. The yeast protein expression system serve as a eukaryotic system integrate the advantages of the mammalian cell expression system. A protein expressed by yeast system could be modificated such as glycosylation, acylation, phosphorylation and so on to ensure the native protein conformation. It can be used to produce protein material with high added value that is very close to the natural protein. Our proteins produced by yeast expression system has been used as raw materials for downstream preparation of monoclonal antibodies.

Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Concentration:	0.2-2 mg/mL
Buffer:	Tris-based buffer, 50 % glycerol
Handling Advice:	Repeated freezing and thawing is not recommended. Store working aliquots at 4 °C for up to one week
Storage:	-20 °C
Storage Comment:	Store at -20 °C, for extended storage, conserve at -20 °C or -80 °C.