

Datasheet for ABIN1672858

Endothelin 1 ELISA Kit

1 Image

2

Publications



Overview

Quantity:	96 tests
Target:	Endothelin 1 (EDN1)
Binding Specificity:	AA 53-73
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	3.9-250 pg/mL
Minimum Detection Limit:	3.9 pg/mL
Application:	ELISA

Product Details

Purpose:	Sandwich High Sensitivity ELISA kit for Quantitative Detection of Mouse Endothelin
Brand:	PicoKine™
Sample Type:	Cell Culture Supernatant, Serum, Plasma (heparin), Plasma (EDTA)
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Immunogen:	Expression system for standard: human peptide Immunogen sequence: C53-W73
Specificity:	Expression system for standard: human peptide Immunogen sequence: C53-W73
Cross-Reactivity (Details):	There is no detectable cross-reactivity with other relevant proteins.

Product Details

Predicted Reactivity:	Hamster
Sensitivity:	<0.5pg/mL
Material not included:	Microplate reader in standard size. Automated plate washer. Adjustable pipettes and pipette tips. Multichannel pipettes are recommended in the condition of large amount of samples in the detection. Clean tubes and Eppendorf tubes. Washing buffer (neutral PBS or TBS). Preparation of 0.01M TBS: Add 1.2g Tris, 8.5g Nacl
Target Details	
Target:	Endothelin 1 (EDN1)
Alternative Name:	EDN1 (EDN1 Products)
Background:	Protein Function: Endothelins are endothelium-derived vasoconstrictor peptides. Background: The endothelins is a family of structurally and pharmacologically distinct peptides, which has been identified and sequenced in humans. Three isoforms of human endothelin have been identified: endothelin-1, -2, and -3. Endothelin-1 is a potent, 21-amino acid vasoconstrictor peptide produced by vascular endothelial cells. Endothelins are 21-amino acid vasoconstricting peptides produced primarily in the endothelium having a key role in vascular homeostasis. Endothelins are implicated in vascular diseases of several organ systems, including the heart, general circulation and brain. Endothelins are proteins that constrict blood vessels and raise blood pressure. They are normally kept in balance by other mechanisms, but when they are over-expressed, they contribute to high blood pressure(hypertension) and heart disease. Synonyms: Endothelin-1,ET-1,Preproendothelin-1,PPET1,Edn1, Full Gene Name: Endothelin-1 Cellular Localisation: Secreted.
Gene ID:	13614
UniProt:	P22387
Pathways:	Hormone Transport, Negative Regulation of Hormone Secretion, Regulation of Systemic Arterial Blood Pressure by Hormones, cAMP Metabolic Process, Regulation of Muscle Cell Differentiation, Regulation of G-Protein Coupled Receptor Protein Signaling, Regulation of Cell Size

Application Details

Application Notes: Before using Kit, spin tubes and bring down all components to bottom of tube. Duplicate well

Application Details

	assay was recommended for both standard and sample testing.
Comment:	Tissue Specificity: Highest expression in the adult is in lung. Lower levels found in heart, kidney
	brain and intestine. In the embryo, expressed in outer and inner pharyngeal arch surfaces. Also
	expressed in endothelium of dorsal aorta and arch arteries, and in epithelium of pharyngeal
	pouches
Plate:	Pre-coated
Protocol:	mouse Endothelin ELISA Kit is based on standard sandwich enzyme-linked immune-sorbent
	assay technology. A monoclonal antibody from rat specific for Endothelin has been precoated
	onto 96-well plates. Standards (mouse peptide, C53-W73) and test samples are added to the
	wells, a biotinylated detection polyclonal antibody from goat specific for Endothelin is added
	subsequently and then followed by washing with PBS or TBS buffer. Avidin-Biotin-Peroxidase
	Complex is added and unbound conjugates are washed away with PBS or TBS buffer. HRP
	substrate TMB are used to visualize HRP enzymatic reaction. TMB is catalyzed by HRP to
	produce a blue color product that changed into yellow after adding acidic stop solution. The
	density of yellow is proportional to the mouse Endothelin amount of sample captured in plate.
Assay Procedure:	Aliquot 0.1 mL per well of the 250pg/mL, 125pg/mL, 62.5pg/mL, 31.2pg/mL, 16.5pg/mL,
	7.8pg/mL, 3.9pg/mL mouse Endothelin standard solutions into the precoated 96-well plate. Ad
	0.1 mL of the sample diluent buffer into the control well (Zero well). Add 0.1 mL of each
	properly diluted sample of mouse cell culture supernatants, serum or plasma(heparin, EDTA) to
	each empty well. See "Sample Dilution Guideline" above for details. We recommend that each
	mouse Endothelin standard solution and each sample is measured in duplicate.
Assay Precision:	• Sample 1: n=16, Mean(pg/ml): 22.2, Standard deviation: 0.777, CV(%): 3.5
	• Sample 2: n=16, Mean(pg/ml): 85.7, Standard deviation: 4.11, CV(%): 4.8
	• Sample 3: n=16, Mean(pg/ml): 144, Standard deviation: 8.21, CV(%): 5.7,
	 Sample 1: n=24, Mean(pg/ml): 21.4, Standard deviation: 1.46, CV(%): 6.8 Sample 2: n=24, Mean(pg/ml): 91.5, Standard deviation: 6.68, CV(%): 7.3
	• Sample 3: n=24, Mean(pg/ml): 153, Standard deviation: 12.09, CV(%): 7.9
Restrictions:	For Research Use only
Handling	
Handling Advice:	Avoid multiple freeze-thaw cycles.
Storage:	-20 °C,4 °C
Storage Comment:	Store at 4°C for 6 months, at -20°C for 12 months. Avoid multiple freeze-thaw cycles

Expiry Date:

12 months


Publications

Product cited in:

Liu, Zhu, Zhou, Wei, Long, Chen, Ling, Ge, Zhuo: "Endoplasmic reticulum stress promotes amyloid-beta peptides production in RGC-5 cells." in: **Cell stress & chaperones**, Vol. 19, Issue 6, pp. 827-35, (2014) (PubMed).

Peng, Dai, Ji, Dai: "The separate roles of endothelin receptors participate in remodeling of matrix metalloproteinase and connexin 43 of cardiac fibroblasts in maladaptive response to isoproterenol." in: **European journal of pharmacology**, Vol. 634, Issue 1-3, pp. 101-6, (2010) (PubMed).

Images



ELISA

Image 1. Mouse Endothelin PicoKine ELISA Kit standard curve