

[Go to Product page](#)

Datasheet for ABIN1674225

Retinoic Acid Induced 12 (RAI12) (AA 1-318) protein (His tag)

Overview

Quantity:	1 mg
Target:	Retinoic Acid Induced 12 (RAI12)
Protein Characteristics:	AA 1-318
Origin:	Rat
Source:	Yeast
Protein Type:	Recombinant
Purification tag / Conjugate:	His tag
Application:	ELISA

Product Details

Sequence:	MPRLSGDDI RRRPEASEML DSSLATGGLV LLRDSVEWEG RGLLKALIKK SALRGEQVHV LGCEVSEEEF REGLGSDVNS RLVYHDLFRD PLNWSQPGEA APEGPLKALR SMCRRTRDRGS VTIALDSLW LLCHIPCVTL CQALHALSQR NVDPGDNPLI EQVRVLGLLH EELHGPGPVG AVSSLAHTEV TLGKMDQTS ASILCRPPQQ RATYQTTWWFS ILPDFSLDLH EGLPLHSELH RDPHTTQVDP ASHLTFNLHL SKKEREAKDS LTLPFQFSSE KQQALLHPVP GQTTGRIFYE PDAFDDVDQE DPDDDLDI
Specificity:	Rattus norvegicus (Rat)
Characteristics:	Please inquire if you are interested in this recombinant protein expressed in E. coli, mammalian cells or by baculovirus infection. Be aware about differences in price and lead time.
Purity:	> 90 %

Target Details

Target:	Retinoic Acid Induced 12 (RAI12)
Alternative Name:	Dermal papilla-derived protein 6 homolog (Derp6) (RAI12 Products)
Background:	Recommended name: Dermal papilla-derived protein 6 homolog
UniProt:	Q6IUP3

Application Details

Comment:	The yeast protein expression system is the most economical and efficient eukaryotic system for secretion and intracellular expression. A protein expressed by the mammalian cell system is of very high-quality and close to the natural protein. But the low expression level, the high cost of medium and the culture conditions restrict the promotion of mammalian cell expression systems. The yeast protein expression system serve as a eukaryotic system integrate the advantages of the mammalian cell expression system. A protein expressed by yeast system could be modified such as glycosylation, acylation, phosphorylation and so on to ensure the native protein conformation. It can be used to produce protein material with high added value that is very close to the natural protein. Our proteins produced by yeast expression system has been used as raw materials for downstream preparation of monoclonal antibodies.
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Concentration:	0.2-2 mg/mL
Buffer:	Tris-based buffer, 50 % glycerol
Handling Advice:	Repeated freezing and thawing is not recommended. Store working aliquots at 4 °C for up to one week
Storage:	-20 °C
Storage Comment:	Store at -20 °C, for extended storage, conserve at -20 °C or -80 °C.