# antibodies - online.com





## NAGK Protein (AA 1-302) (His tag)



$\sim$					
( )	\/	Δ	r١	/1	۱۸

Characteristics:

> 90 %

Purity:

Quantity:	1 mg
Target:	NAGK
Protein Characteristics:	AA 1-302
Origin:	Vibrio parahaemolyticus
Source:	Yeast
Protein Type:	Recombinant
Purification tag / Conjugate:	This NAGK protein is labelled with His tag.
Application:	ELISA
Product Details	
Sequence:	MYYGFDVGGT KIEFGAFNEK LERVATERVP TPTDNYELLV DTIAELVNKY DAEFGCEGTI
	GLGLPGMEDA DDATVLTVNV PAAKGKPLRA DLEAKIGRSV KIENDANCFA LSEAWDEELQ
	DEPSVLGLIL GTGFGGGFIY DGKVFSGRNH VAGEVGHTRL PIDAWFHLGE NAPLLGCGCD
	KKGCLDSYLS GRGFELLYAH YYGEEKKAID IIKAHAEGEA KAVEHVERFM ELLAICFANI
	FTATDPHVVV LGGGLSNFEL IYEEMPKRIP KYLLSVAKCP KIIKAKHGDS GGVRGAAFLH IK
Specificity:	Vibrio parahaemolyticus

Please inquire if you are interested in this recombinant protein expressed in E. coli, mammalien

cells or by baculovirus infection. Be aware about differences in price and lead time.

#### **Target Details**

Target:	NAGK	
Alternative Name:	N-acetyl-D-glucosamine kinase (nagK) (NAGK Products)	
Background:	Recommended name: N-acetyl-D-glucosamine kinase.  EC= 2.7.1.59.  Alternative name(s): GlcNAc kinase	
UniProt:	Q87PK8	

#### **Application Details**

#### Comment:

The yeast protein expression system is the most economical and efficient eukaryotic system for secretion and intracellular expression. A protein expressed by the mammalian cell system is of very high-quality and close to the natural protein. But the low expression level, the high cost of medium and the culture conditions restrict the promotion of mammalian cell expression systems. The yeast protein expression system serve as a eukaryotic system integrate the advantages of the mammalian cell expression system. A protein expressed by yeast system could be modificated such as glycosylation, acylation, phosphorylation and so on to ensure the native protein conformation. It can be used to produce protein material with high added value that is very close to the natural protein. Our proteins produced by yeast expression system has been used as raw materials for downstream preparation of monoclonal antibodies.

Restrictions:

For Research Use only

### Handling

Format:	Lyophilized
Concentration:	0.2-2 mg/mL
Buffer:	Tris-based buffer, 50 % glycerol
Handling Advice:	Repeated freezing and thawing is not recommended. Store working aliquots at 4 °C for up to one week
Storage:	-20 °C
Storage Comment:	Store at -20 °C, for extended storage, conserve at -20 °C or -80 °C.