

Datasheet for ABIN1684628 IL17RC Protein (C-Term, Extracellular Domain)



Overview

000101000	
Quantity:	100 µg
Target:	IL17RC
Protein Characteristics:	C-Term, Extracellular Domain
Origin:	Human
Source:	HEK-293T Cells
Protein Type:	Recombinant
Product Details	
Specificity:	Optimized DNA sequence encoding extracellular domain of human IL-17RC receptor including a C terminal Human IgG1 Fc tag was expressed in HEK293 cells.
Characteristics:	Recombinant IL17RC-Fc is a homodimer protein consisting of 675 amino acid residue subunits, due to glycosylation migrates as an approximately 100kD protein under reducing SDS-PAGE conditions.
Purity:	> 98 %, as determined by SDS-PAGE and HPLC
Sterility:	0.2 µm filtered
Endotoxin Level:	Endotoxin content was assayed using a LAL gel clot method. Endotoxin level was found to be less than 0.1 ng/µg(1EU/µg).
Target Details	

Target:	IL17RC
Alternative Name:	IL17RC (IL17RC Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1684628 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Background:	Common gamma chain Interleukin (IL)-2, IL-4, IL-7, IL-9 and IL-15 belong to a family of cytokines whose receptors share the common gamma chain (gamma c). Despite some redundancy, thesecytokines that signal through the gamma c (gamma c cytokines) have essentially distinct and oftencritical effects in normal T-cell development, survival, proliferation and differentiation. Importantly, they have also been implicated, directly or indirectly, in T-cell leukemogenesis.
UniProt:	Q8NAC3
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Buffer:	PBS solution, pH7.2
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	-20 °C
Storage Comment:	The lyophilized protein is stable for at least years from date of receipt at -20 °C. Upon reconstitution, this cytokine can be stored in working aliquots at - 8 °C for one month, or at -20 °C for six months, with a carrier protein without detectable loss of activity.

Expiry Date: 12-24 months