



[Go to Product page](#)

Datasheet for ABIN1684639

Leptin Receptor Protein (LEPR) (C-Term, Extracellular Domain)

Overview

Quantity:	100 µg
Target:	Leptin Receptor (LEPR)
Protein Characteristics:	C-Term, Extracellular Domain
Origin:	Human
Source:	HEK-293T Cells
Protein Type:	Recombinant
Biological Activity:	Active

Product Details

Specificity:	Optimized DNA sequence encoding extracellular domain of human leptin receptor including a C-terminal Human IgG1 Fc tag was expressed in HEK293 cells.
Characteristics:	Recombinant human leptin receptor is a homodimer protein consisting of two 1070 amino acid residue subunit chains, due to glycosylation migrates as an approximately 160kDa protein on SDS-PAGE.
Purity:	> 98 %, as determined by SDS-PAGE and HPLC
Sterility:	0.2 µm filtered
Endotoxin Level:	Endotoxin content was assayed using a LAL gel clot method. Endotoxin level was found to be less than 0.1 ng/µg(1EU/µg).

Target Details

Target:	Leptin Receptor (LEPR)
---------	------------------------

Target Details

Alternative Name: [LEPR \(LEPR Products\)](#)

Background: FGF-10 is involved in the initial budding as well as the continuous outgrowth of vertebrate limbs, FGF10 mRNA is expressed preferentially in neurons but not in glial cells and may have a distinct role in the brain. Human FGF-10 is mitogenic for fetal rat keratinizing epidermal cells but not for NIH 3T3 cells. Recombinant FGF-10 induces the proliferation of human urothelial cells in vitro and induces the proliferation of transitional epithelium. FGF-10 is secreted by cultured mouse pre-adipocytes, prevention of FGF-10 signaling inhibits subsequent differentiation. The ability of embryonic fibroblasts derived from FGF-10 knock-out mice to differentiate into adipocytes is also impaired.

UniProt: [P48357](#)

Pathways: [JAK-STAT Signaling](#), [AMPK Signaling](#), [Feeding Behaviour](#)

Application Details

Comment: Biologically active: Measured by its ability to bind immobilized recombinant human Leptin (1 µg/mL) at a linear concentration range of 0.01-4 µg/mL.

Restrictions: For Research Use only

Handling

Format: Lyophilized

Buffer: PBS solution, pH7.2

Handling Advice: Avoid repeated freeze/thaw cycles.

Storage: -20 °C

Storage Comment: The lyophilized antibody is stable for at least 1 year from date of receipt at -20 °C. Upon reconstitution, this antibody can be stored in working aliquots at -8 °C for one month, or at -20 °C for six months without detectable loss of activity.

Expiry Date: 12 months
