
# antibodies -online.com





# anti-TNFRSF1A antibody (AA 20-43)



**Images** 



Go to Product page

$\sim$			
	$  \backslash / \cap$	r\/I	$\triangle V$

Quantity:	100 μg	
Target:	TNFRSF1A	
Binding Specificity:	AA 20-43	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TNFRSF1A antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunofluorescence (IF), Immunocytochemistry (ICC)	
Product Details		
Immunogen:	Peptide corresponding to AA 20-43 of the mouse TNF-R1 sequence, identical to rat and human over those residues	
Specificity:	Detects $\sim$ 55 kDa. Other bands present may be the result of oligomerization, self-aggregation and/or cleavage of the TNF-R1 extracellular domain.	
Cross-Reactivity:	Cow, Dog, Human, Monkey, Mouse, Rabbit, Rat	
Purification:	Peptide Affinity Purified	
Target Details		
Target:	TNFRSF1A	
Alternative Name:	TNFR1 (TNFRSF1A Products)	


## Target Details


rarget betails		
Background:	The Tumor Necrosis Factor Receptor (TNFR) also known as Cluster of differentiation (CD120) is a protein that belongs to the (TNF)/ (TNFR) superfamily. TNF interacts with two distinct receptors TNFR1 and TNFR2. These receptors share no homology on their cytoplasmic sequences(1,3).TNFR1 also known as p55/p60 is a high affinity receptor for TNF-a. The TNFR1 has an extracellular domain with variable numbers of cysteine-rich repeats. The functional properties of TNFR1 are targets in new therapies for osteoporosis, chronic inflammatory and autoimmune diseases (1, 2). The TNF-a/TNFR1 receptor complex is responsible for the recruitment and the subsequent activation of the caspase (aspartate-specific cysteine proteases) that regulate apoptosis.	
Gene ID:	8666	
UniProt:	P19438	
Pathways:	NF-kappaB Signaling, Apoptosis, Caspase Cascade in Apoptosis, Hepatitis C, Ubiquitin Proteasome Pathway	
Application Details		
Application Notes:	<ul> <li>WB (1:1000)</li> <li>IHC (1:100)</li> <li>ICC/IF (1:100)</li> <li>optimal dilutions for assays should be determined by the user.</li> </ul>	
Comment:	1 μg/ml of ABIN1686656 was sufficient for detection of TNFR1 in 20 μg of Hela lysate by colorimetric immunoblot analysis using Goat anti-rabbit IgG:HRP as the secondary antibody.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C	

Storage Comment:

-20°C

#### **Images**






#### **Western Blotting**

Image 1. Western blot analysis of Human A549 showing detection of ~ 50 kDa TNF-R1 protein using Rabbit Anti-TNF-R1 Polyclonal Antibody (ABIN1686656). Lane 1: MW Ladder, Lane 2: A549. Load: 30 µg. Block: 5 % BSA in TBST. Primary Antibody: Rabbit Anti-TNF-R1 Polyclonal Antibody (ABIN1686656) at 1:1000 for 2 hours at RT with shaking. Secondary Antibody: Goat Anti-Rabbit IgG: HRP at 1:4000 for 1 hour at RT with shaking. Color Development: Chemiluminescent for HRP (Moss) for 5 min in RT. Predicted/Observed Size: ~ 50 kDa. Other Band(s): ~90-100 kDa. Other bands can be explained by a few factors, such as oligomerization, self-aggregation, cleavage of the TNFR1 extracellular domain, etc.( Literature references: doi: 10.3389/fcell.2020.615141, 10.1128/MCB.22.8.2536-2543.2002, 10.1073/pnas.0307981100).

#### **Western Blotting**

Image 2. Western blot analysis of Mouse Liver cell lysates showing detection of  $\sim 55$  kDa TNF-R1 protein using Rabbit Anti-TNF-R1 Polyclonal Antibody . Lane 1: Molecular Weight Ladder (MW). Lane 2: Mouse Liver cell lysates. Load: 15 µg. Block: 5% Skim Milk in 1X TBST. Primary Antibody: Rabbit Anti-TNF-R1 Polyclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Goat Anti-Rabbit IgG: HRP at 1:2000 for 60 min at RT. Color Development: ECL solution for 5 min at RT. Predicted/Observed Size:  $\sim 55$  kDa.



### **Immunohistochemistry**

Image 3. Immunohistochemistry analysis using Rabbit Anti-TNF-R1 Polyclonal Antibody . Tissue: backskin. Species: Mouse. Fixation: Bouin's Fixative Solution. Primary Antibody: Rabbit Anti-TNF-R1 Polyclonal Antibody at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Rabbit (green) at 1:50 for 1 hour at RT. Localization: dermis.

Please check the product details page for more images. Overall 6 images are available for ABIN1686656.