

Datasheet for ABIN1691578

Cathepsin L Protein (AA 18-334) (His tag)



Overview

Quantity:	50 μg	
Target:	Cathepsin L (CTSL1)	
Protein Characteristics:	AA 18-334	
Origin:	Mouse	
Source:	Human Cells	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This Cathepsin L protein is labelled with His tag.	

Product Details

Product Details			
Purpose:	Recombinant Mouse Cathepsin L/CTSL (C-6His)		
Sequence:	TPKFDQTFSA EWHQWKSTHR RLYGTNEEEW RRAIWEKNMR MIQLHNGEYS NGQHGFSMEM		
	NAFGDMTNEE FRQVVNGYRH QKHKKGRLFQ EPLMLKIPKS VDWREKGCVT PVKNQGQCGS		
	CWAFSASGCL EGQMFLKTGK LISLSEQNLV DCSHAQGNQG CNGGLMDFAF QYIKENGGLD		
	SEESYPYEAK DGSCKYRAEF AVANDTGFVD IPQQEKALMK AVATVGPISV AMDASHPSLQ		
	FYSSGIYYEP NCSSKNLDHG VLLVGYGYEG TDSNKNKYWL VKNSWGSEWG MEGYIKIAKD		
	RDNHCGLATA ASYPVVNVDH HHHHH		
Characteristics:	Recombinant Mouse Cathepsin L/Ctsl1 is produced by our mammalian expression system in		
	human cells. The target protein is expressed with sequence (Thr18-Asn334) of Mouse		
	Cathepsin L fused with a polyhistidine tag at the C-terminus.		
Purity:	> 95 % as determined by reducing SDS-PAGE.		
Sterility:	0.2 μm filtered		

Target Details Target: Cathepsin L (CTSL1) Alternative Name: Cathepsin L (CTSL1 Products) Sub Type: Fusionprotein Background: Mouse Cathepsin L is a lysosomal cysteine proteinase which is a member of the peptidase C1 family. This protein is a dimer composed of disulfide-linked heavy and light chains, both produced from a single protein precursor. Cathepsin L plays a major role in intracellular protein catabolism. Its substrates include collagen and elastin, as well as alpha-1 protease inhibitor, a major controlling element of neutrophil elastase activity. Cathepsin L has been implicated in several pathologic processes, including myofforil necrosis in myopathies and in myocardial ischemia, and in the renal tubular response to proteinuria. Alternative Names: Cathepsin L 1, Major excreted protein, p39 cysteine proteinase, Cts11, Molecular Weight: 36.8 kDa UniProt: P06797 Pathways: Activation of innate immune Response, Toll-Like Receptors Cascades Application Details Restrictions: For Research Use only Handling Format: Lyophilized It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Buffer: Lyophilized from a 0.2 µm filtered solution of 20 mM PB,150 mM NaCl, pH 7.4. Handling Advice: Always centrifuge tubes before opening. Do not mix by vortex or pipetting.	Product Details				
Alternative Name: Cathepsin L (CTSL1) Alternative Name: Cathepsin L (CTSL1 Products) Sub Type: Fusionprotein Mouse Cathepsin L is a lysosomal cysteine proteinase which is a member of the peptidase C1 family. This protein is a dimer composed of disulfide-linked heavy and light chains, both produced from a single protein precursor. Cathepsin L plays a major role in intracellular proteir catabolism. Its substrates include collagen and elastin, as well as alpha-1 protease inhibitor, a major controlling element of neutrophil elastase activity. Cathepsin L has been implicated in several pathologic processes, including myofibrin necrosis in myopathies and in myocardial ischemia, and in the renal tubular response to proteinuria. Alternative Names: Cathepsin L1, Major excreted protein, p39 cysteine proteinase, Cts11, Molecular Weight: 36.8 kDa UniProt: P06797 Pathways: Activation of Innate Immune Response, Toll-Like Receptors Cascades Application Details Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles Buffer: Lyophilized from a 0.2 µm filtered solution of 20 mM P8,150 mM NaCl, pH 7.4. Handling Advice: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Storage: 4 *CC-20 *CC-80 *CC Storage Comment: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks	Endotoxin Level:	Less than 0.1 ng/μg (1 IEU/μg) as determined by LAL test			
Alternative Name: Cathepsin L (CTSL1 Products) Sub Type: Fusionprotein Mouse Cathepsin L is a lysosomal cysteine proteinase which is a member of the peptidase C1 family. This protein is a dimer composed of disulfide-linked heavy and light chains, both produced from a single protein precursor. Cathepsin L plays a major role in intracellular proteir catabolism. Its substrates include collagen and elastin, as well as alpha-1 protease inhibitor, a major controlling element of neutrophil elastase activity. Cathepsin L has been implicated in several pathologic processes, including myofibril necrosis in myopathies and in myocardial ischemia, and in the renal tubular response to proteinuria. Alternative Names: Cathepsin L1, Major excreted protein, p39 cysteine proteinase, Ctsl1, Molecular Weight: 36.8 kDa UniProt: P06797 Pathways: Activation of Innate Immune Response, Toll-Like Receptors Cascades Application Details Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Buffer: Lyophilized from a 0.2 µm filtered solution of 20 mM P8,150 mM NaCl, pH 7.4. Handling Advice: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Storage: 4 "C/-20 "C/-80 "C Storage Comment: Lyophilized protein should be stored at <-20°C, though stable at room temperature for 3 weeks	Target Details				
Sub Type: Fusionprotein Background: Mouse Cathepsin L is a lysosomal cysteine proteinase which is a member of the peptidase C1 family. This protein is a dimer composed of disulfide-linked heavy and light chains, both produced from a single protein precursor. Cathepsin L plays a major role in intracellular protein catabolism. Its substrates include collagen and elastin, as well as alpha-1 protease inhibitor, a major controlling element of neutrophil elastase activity. Cathepsin L has been implicated in several pathologic processes, including myofibril necrosis in myopathies and in myocardial ischemia, and in the renal tubular response to proteinuria. Alternative Names: Cathepsin L 1, Major excreted protein, p39 cysteine proteinase, Ctsl1, Molecular Weight: 36.8 kDa UniProt: P06797 Pathways: Activation of Innate immune Response, Toll-Like Receptors Cascades Application Details Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in ddH20. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Buffer: Lyophilized from a 0.2 µm filtered solution of 20 mM PB,150 mM NaCl, pH 7.4. Handling Advice: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Storage: 4 °C/-20 °C/-80 °C Storage Comment: Lyophilized protein should be stored at <-20°C, though stable at room temperature for 3 weeks	Target:	Cathepsin L (CTSL1)			
Background: Mouse Cathepsin L is a lysosomal cysteine proteinase which is a member of the peptidase C1 family. This protein is a dimer composed of disulfide-linked heavy and light chains, both produced from a single protein precursor. Cathepsin L plays a major role in intracellular protein catabolism. Its substrates include collagen and elastin, as well as alpha-1 protease inhibitor, a major controlling element of neutrophile lastase activity. Cathepsin L has been implicated in several pathologic processes, including myofibril necrosis in myopathies and in myocardial ischemia, and in the renal tubular response to proteinuria. Alternative Names: Cathepsin L1, Major excreted protein, p39 cysteine proteinase, Ctsl1, Molecular Weight: 36.8 kDa UniProt: P06797 Pathways: Activation of Innate immune Response, Toll-Like Receptors Cascades Application Details Restrictions: For Research Use only Handling Format: Lyophilized It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in ddH20. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Buffer: Lyophilized from a 0.2 µm filtered solution of 20 mM PB,150 mM NaCl, pH 7.4. Handling Advice: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Storage: 4 *C/-20 *C/-80 *C Storage Comment: Lyophilized protein should be stored at < -20 *C, though stable at room temperature for 3 weeks	Alternative Name:	Cathepsin L (CTSL1 Products)			
family. This protein is a dimer composed of disulfide-linked heavy and light chains, both produced from a single protein precursor. Cathepsin L plays a major role in intracellular proteir catabolism. Its substrates include collagen and elastin, as well as alpha-1 protease inhibitor, a major controlling element of neutrophil elastase activity. Cathepsin L has been implicated in several pathologic processes, including myofibril necrosis in myopathies and in myocardial ischemia, and in the renal tubular response to proteinuria. Alternative Names: Cathepsin L1, Major excreted protein, p39 cysteine proteinase, Ctsl1, Molecular Weight: 906797 Pathways: Activation of Innate immune Response, Toll-Like Receptors Cascades Application Details Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in ddH20. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Buffer: Lyophilized from a 0.2 µm filtered solution of 20 mM PB,150 mM NaCl, pH 7.4. Handling Advice: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Storage: 4 *C/-20 *C/-80 *C Storage Comment: Lyophilized protein should be stored at <-20*C, though stable at room temperature for 3 weeks	Sub Type:	Fusionprotein			
UniProt: P06797 Pathways: Activation of Innate immune Response, Toll-Like Receptors Cascades Application Details Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in ddH20. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Buffer: Lyophilized from a 0.2 µm filtered solution of 20 mM PB,150 mM NaCl, pH 7.4. Handling Advice: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Storage: 4 °C/-20 °C/-80 °C Storage Comment: Lyophilized protein should be stored at < -20 °C, though stable at room temperature for 3 weeks	Background:	family. This protein is a dimer composed of disulfide-linked heavy and light chains, both produced from a single protein precursor. Cathepsin L plays a major role in intracellular protein catabolism. Its substrates include collagen and elastin, as well as alpha-1 protease inhibitor, a major controlling element of neutrophil elastase activity. Cathepsin L has been implicated in several pathologic processes, including myofibril necrosis in myopathies and in myocardial ischemia, and in the renal tubular response to proteinuria.			
Pathways: Activation of Innate immune Response, Toll-Like Receptors Cascades Application Details Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Buffer: Lyophilized from a 0.2 µm filtered solution of 20 mM PB,150 mM NaCl, pH 7.4. Handling Advice: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Storage: 4 °C/-20 °C/-80 °C Storage Comment: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks	Molecular Weight:	36.8 kDa			
Application Details Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Buffer: Lyophilized from a 0.2 µm filtered solution of 20 mM PB,150 mM NaCl, pH 7.4. Handling Advice: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Storage: 4 °C/-20 °C/-80 °C Storage Comment: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks	UniProt:	P06797			
For Research Use only Handling Format: Lyophilized Reconstitution: It is not recommended to reconstitute to a concentration less than 100 μg/mL. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Buffer: Lyophilized from a 0.2 μm filtered solution of 20 mM PB,150 mM NaCl, pH 7.4. Handling Advice: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Storage: 4 °C/-20 °C/-80 °C Storage Comment: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks	Pathways:	Activation of Innate immune Response, Toll-Like Receptors Cascades			
Handling Format: Lyophilized Reconstitution: It is not recommended to reconstitute to a concentration less than 100 μg/mL. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Buffer: Lyophilized from a 0.2 μm filtered solution of 20 mM PB,150 mM NaCl, pH 7.4. Handling Advice: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Storage: 4 °C/-20 °C/-80 °C Storage Comment: Lyophilized protein should be stored at < -20 °C, though stable at room temperature for 3 weeks	Application Details				
Format: Lyophilized Reconstitution: It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in ddH20. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Buffer: Lyophilized from a 0.2 µm filtered solution of 20 mM PB,150 mM NaCl, pH 7.4. Handling Advice: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Storage: 4 °C/-20 °C/-80 °C Storage Comment: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks	Restrictions:	For Research Use only			
Reconstitution: It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Buffer: Lyophilized from a 0.2 µm filtered solution of 20 mM PB,150 mM NaCl, pH 7.4. Handling Advice: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Storage: 4 °C/-20 °C/-80 °C Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks	Handling				
Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Buffer: Lyophilized from a 0.2 µm filtered solution of 20 mM PB,150 mM NaCl, pH 7.4. Handling Advice: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Storage: 4 °C/-20 °C/-80 °C Lyophilized protein should be stored at < -20 °C, though stable at room temperature for 3 weeks	Format:	Lyophilized			
Handling Advice: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Storage: 4 °C/-20 °C/-80 °C Lyophilized protein should be stored at < -20 °C, though stable at room temperature for 3 weeks	Reconstitution:	Dissolve the lyophilized protein in ddH2O.			
Storage: 4 °C/-20 °C/-80 °C Storage Comment: Lyophilized protein should be stored at < -20 °C, though stable at room temperature for 3 weeks	Buffer:	Lyophilized from a 0.2 µm filtered solution of 20 mM PB,150 mM NaCl, pH 7.4.			
Storage Comment: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks	Handling Advice:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting.			
	Storage:	4 °C/-20 °C/-80 °C			
	Storage Comment:	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks Reconstituted protein solution can be stored at 4-7°C for 2-7 days.			

1	-	0	ИI	0	_
$\overline{}$	171	11(111	11	(

Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Expiry Date: 6 months