# antibodies .- online.com





# anti-C6ORF195 antibody (AA 81-127) (Alexa Fluor 350)



Go to Product page

| ( ) | 1/0 | r\ /1 | 014 |   |
|-----|-----|-------|-----|---|
| ( ) | ve  | I V I | -v  | V |
|     |     |       |     |   |

| Quantity:            | 100 μL  |  |
|----------------------|---|--|
| Target:              | C60RF195  |  |
| Binding Specificity: | AA 81-127   |  |
| Reactivity:          | Human   |  |
| Host:                | Rabbit  |  |
| Clonality:           | Polyclonal  |  |
| Conjugate:           | This C6ORF195 antibody is conjugated to Alexa Fluor 350   |  |
| Application:         | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |  |

## **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human C6orf195 |  |
|-----------------------|--|--|
| Isotype:              | IgG  |  |
| Predicted Reactivity: | d Reactivity: Human  |  |
| Purification:         | Purified by Protein A.                                       |  |

# **Target Details**

| Target:           | C60RF195   |  |
|-------------------|--|--|
| Alternative Name: | C6orf195 (C6ORF195 Products)   |  |
| Background:       | Synonyms: bA145H9.2, C6orf195, CF195_HUMAN, Chromosome 6 open reading frame 195, |  |

FLJ31934, RP11-145H9.2, Uncharacterized protein C6orf195.

Background: Making up nearly 6 % of the human genome, chromosome 6 contains around 1,200 genes within 170 million base pairs of sequence. Deletion of a portion of the q arm of chromosome 6 is associated with early onset intestinal cancer suggesting the presence of a cancer susceptibility locus. Porphyria cutanea tarda is associated with chromosome 6 through the HFE gene which, when mutated, predisposes an individual to developing this porphyria. Notably, the PARK2 gene, which is associated with Parkinson's disease, and the genes encoding the major histocompatiblity complex proteins, which are key molecular components of the immune system and determine predisposition to rheumatic diseases, are also located on chromosome 6. Stickler syndrome, 21-hydroxylase deficiency and maple syrup urine disease are also associated with genes on chromosome 6. A bipolar disorder susceptibility locus has been identified on the q arm of chromosome 6. The C6orf195 gene product has been provisionally designated C6orf195 pending further characterization.

Gene ID:

154386

### **Application Details**

| Application Notes: |  |
|--------------------|--|

IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months  |