

[Go to Product page](#)

Datasheet for ABIN1692724

anti-FANCM antibody (AA 831-930) (Alexa Fluor 350)

Overview

Quantity:	100 µL
Target:	FANCM
Binding Specificity:	AA 831-930
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FANCM antibody is conjugated to Alexa Fluor 350
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FANCM
Isotype:	IgG
Predicted Reactivity:	Human
Purification:	Purified by Protein A.

Target Details

Target:	FANCM
Alternative Name:	FANCM (FANCM Products)
Background:	Synonyms: FAAP250, Fanconi anemia group M protein, Protein Hef ortholog,

Target Details

Background: Fanconi anemia (FA) is an autosomal recessive disorder characterized by bone marrow failure, birth defects and chromosomal instability. At the cellular level, FA is characterized by spontaneous chromosomal breakage and a unique hypersensitivity to DNA cross-linking agents. The thirteen FA proteins that have been characterized are important for regulating chromosomal stability and genome surveillance. Eight of these proteins, namely FANCA, FANCB, FANCC, FANCE, FANCF, FANCG, FANCL and FANCM, comprise the FA core complex, which catalyzes a key reaction in DNA repair: the monoubiquitination of FANCD2. FANCM (Fanconi anemia, complementation group M) is a member of the DEAD-box helicase family of proteins and contains a DEAH helicase domain and a nuclease domain. Localizing to chromatin fractions, FANCM is phosphorylated in a cell cycle-dependent manner and is believed to function as an anchor, recruiting the FA core complex to chromatin. Mutations in the gene encoding FANCM can lead to Fanconi anemia.

Gene ID: 57697

Pathways: [DNA Damage Repair](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months