antibodies -online.com





anti-HIBADH antibody (AA 251-336) (Alexa Fluor 350)



Go to Product page

| () | 11/0 | K\ / | iew | 1 |
|-----|--------|--------|---------------------|---|
| | \cup | 'I V/I | $\square \vee \vee$ | ı |
| | | | | |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | HIBADH |
| Binding Specificity: | AA 251-336 |
| Reactivity: | Human, Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This HIBADH antibody is conjugated to Alexa Fluor 350 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human HIBADH | |
|-----------------------|--|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Human, Mouse | |
| Predicted Reactivity: | Rat,Cow,Sheep,Horse | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | HIBADH |
|-------------------|--------------------------|
| Alternative Name: | Hibadh (HIBADH Products) |

Target Details

Storage Comment:

Expiry Date:

12 months

| rargorbotano | | |
|---------------------|--|--|
| Background: | Synonyms: 3 hydroxy 2 methylpropanoate:NAD+ oxidoreductase, 3 hydroxyisobutyrate | |
| | dehydrogenase, 3 hydroxyisobutyrate dehydrogenase mitochondrial, EC 1.1.1.31, MGC40361, | |
| | NS5ATP1, 3HIDH_HUMAN. | |
| | Background: HIBADH is a 336 amino acid mitochondrial enzyme that catalyzes the NAD+- | |
| | dependent, reversible oxidization of 3-Hydroxyisobutyrate to methylmalonate semialdehyde, an | |
| | intermediate of valine catabolism. The enzyme functions as a homodimer between a pH of 7.0 | |
| | and 10.0, with optimal activity between 8.8 and 9.0. It was previously hypothesized that defects | |
| | in the gene encoding HIBADH may be the cause of 3-Hydroxyisobutyric aciduria, a rare disorder | |
| | that is characterized by a variety of clinical manifestations such as neurodevelopmental | |
| | problems and dysmorphic features. However, it was shown that HIBADH activity was equal in | |
| | patients with 3-Hydroxyisobutyric aciduria as compared with controls. | |
| Gene ID: | 11112 | |
| | | |
| Application Details | | |
| Application Notes: | IF(IHC-P) 1:50-200 | |
| | IF(IHC-F) 1:50-200 | |
| | IF(ICC) 1:50-200 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | |
| | 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | |
| | handled by trained staff only. | |
| Storage: | -20 °C | |
| 04 | Ohanna ah OOOO Alimaah inta madkinla siala ta saasid maasah al faasaa tha saasid | |

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.