# antibodies - online.com





## anti-DOK7 antibody (AA 21-120) (Alexa Fluor 350)



$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:	100 μL	
Target:	DOK7	
Binding Specificity:	AA 21-120	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This DOK7 antibody is conjugated to Alexa Fluor 350	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human DOK7	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Human,Mouse,Dog,Cow,Sheep,Pig,Horse	
Purification:	Purified by Protein A.	

#### **Target Details**

Target:	DOK7
Alternative Name:	DOK7 (DOK7 Products)

### **Target Details**

Background:	Synonyms: Docking protein 7, DOK 7, DOK7, DOK7_HUMAN, Downstream of tyrosine kinase 7,		
	Protein Dok-7.		
	Background: The downstream of kinase family (Dok1-7) are members of a class of docking?		
	proteins that include the tyrosine kinase substrates IRS-1 and Cas, which contain multiple		
	tyrosine residues and putative SH2 binding sites. Based on their similarities, the Dok family of		
	proteins can be divided into three subgroups: Dok-1/2/3, Dok-4/5/6 and Dok-7. Through its		
	interaction with muscle-specific receptor kinase (MuSK), Dok-7 is crucial for neuromuscular		
	synaptogenesis and for MuSK activation. Mice lacking Dok-7 do not form neuromuscular		
	synapses nor acetylcholine receptor clusters. Mutations in the Dok-7 gene can cause		
	congenital myasthenic syndromes (CMA) recessively inherited disorders characterized by		
	muscle weakness.		
Pathways:	Skeletal Muscle Fiber Development		
Application Details			
Application Notes:	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		
Expiry Date:	12 months		