

Datasheet for ABIN1692967

**anti-Cadherin 20 antibody (AA 301-400) (AbBy Fluor® 350)**[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	Cadherin 20 (CDH20)
Binding Specificity:	AA 301-400
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Cadherin 20 antibody is conjugated to AbBy Fluor® 350
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Cadherin 20
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Pig, Horse, Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	Cadherin 20 (CDH20)
Alternative Name:	Cadherin 20 ( <a href="#">CDH20 Products</a> )
Background:	Synonyms: Cadherin 20, cadherin-20, Cadherin 20, type 2, Cdh7, CDH7L3, FLJ37047.

## Target Details

---

Background: The cadherins are a family of Ca<sup>2+</sup>-dependent adhesion molecules that function to mediate cell-cell binding critical to the maintenance of structure and morphogenesis. Cadherins each contain a large extracellular domain at the N-terminus, which is characterized by a series of five homologous repeats, the most distal of which is thought to be responsible for binding specificity. Cadherin-20, also known as protocadherin gamma B4, is a member of the protocadherin gamma gene cluster. Expressed primarily in fibroblasts, cadherin-20 plays a critical role in neuronal cell-cell connections in the brain. Through its ability to mediate cell-cell adhesion, cadherin-20 is thought to help the body respond to injury by regulating post-traumatic wound healing.

## Application Details

---

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months