# antibodies -online.com





## anti-DYNC1I1 antibody (AA 1-100) (Alexa Fluor 350)



Go to Product page

$\sim$	
( )\/白	rview
OVC	

Quantity:	100 μL
Target:	DYNC1I1
Binding Specificity:	AA 1-100
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DYNC1I1 antibody is conjugated to Alexa Fluor 350
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human DYNC1I1
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Cow,Sheep,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

#### Target Details

Target:	DYNC1I1
Alternative Name:	DYNC1I1 (DYNC1I1 Products)

#### **Target Details**

Background:	Synonyms: dynein, cytoplasmic, intermediate polypeptide 1, cytoplasmic dynein 1 intermediate
	chain 1, DH IC-1, dynein intermediate chain 1, cytosolic, cytoplasmic dynein intermediate chain
	1, DC1I1_HUMAN, DYNC1I1, DNCI1, DNCIC1, Dynein cytoplasmic intermediate polypeptide 1.
	Background: This gene is a member of the Tyr protein kinase family and the epidermal growth
	factor receptor subfamily. It encodes a single-pass type I membrane protein with multiple
	cysteine rich domains, a transmembrane domain, a tyrosine kinase domain, a
	phosphotidylinositol-3 kinase binding site and a PDZ domain binding motif. The protein binds to
	and is activated by neuregulins and other factors and induces a variety of cellular responses
	including mitogenesis and differentiation. Multiple proteolytic events allow for the release of a
	cytoplasmic fragment and an extracellular fragment. Mutations in this gene have been
	associated with cancer. Alternatively spliced variants which encode different protein isoforms
	have been described, however, not all variants have been fully characterized. [provided by
	RefSeq, Jul 2008].
Gene ID:	1780
UniProt:	014576
Application Details	
Application Notes:	FCM 1:20-100
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C

### Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months