

[Go to Product page](#)

Datasheet for ABIN1693208

anti-TESK2 antibody (AA 301-400) (Alexa Fluor 350)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | TESK2 |
| Binding Specificity: | AA 301-400 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This TESK2 antibody is conjugated to Alexa Fluor 350 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-----------------------|---|
| Immunogen: | KLH conjugated synthetic peptide derived from human TESK2 |
| Isotype: | IgG |
| Predicted Reactivity: | Human,Mouse,Rat,Horse,Chicken,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|---|
| Target: | TESK2 |
| Alternative Name: | TESK2 (TESK2 Products) |
| Background: | Synonyms: Dual specificity testis specic protein kinase 2, Dual specificity testis-specific protein |

Target Details

kinase 2, MGC29168, OTTHUMP00000009093, RP23-109A3.2, Tesk2, TESK2_HUMAN, Testicular protein kinase 2, Testis specic kinase 2.

Background: TESK2 is a nuclear protein that belongs to the protein kinase superfamily and is expressed in testis and prostate. Functioning as a dual-specificity protein kinase, TESK2 catalyzes the ATP-dependent phosphorylation of substrates and autophosphorylation on tyrosine and serine/threonine residues, thereby mediating intracellular signal transduction pathways. TESK2 requires magnesium as a cofactor and its catalytic activity is thought to play an important role in meiotic events such as spermatogenesis. TESK2 contains one protein kinase domain that is 65 % identical to the kinase domain found in TESK1 (testicular protein kinase 1), suggesting a similar role for these proteins in phosphorylation events. Three isoforms of TESK2 are expressed due to alternative splicing.

Gene ID: 10420

Pathways: [Cell-Cell Junction Organization](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months