## antibodies .- online.com





## anti-NGEF antibody (pTyr179) (AbBy Fluor® 350)



Go to Product page

Overview	
Quantity:	100 μL
Target:	NGEF
Binding Specificity:	pTyr179
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NGEF antibody is conjugated to AbBy Fluor® 350
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human Ephexin-1 around the phosphorylation site of Tyr179
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Chicken
Purification:	Purified by Protein A.
Target Details	
Target:	NGEF

## Target Details

Synonyms: Ephexin-1 phospho Y179, p-Ephexin-1 phospho Y179, ARHGEF27, Eph-interacting exchange protein, EPHEXIN, Ephexin 1, Neuronal guanine nucleotide exchange factor, Ngef, NGEF_HUMAN.  Background: EPHEXIN is a 710 amino acid protein that localizes to both the membrane and the cytoplasm and contains one SH3 domain, one PH domain and one DH domain. Expressed at high levels in brain and present at lower levels in lung tissue, EPHEXIN interacts with EphA4 and functions as a guanine nucleotide exchange factor (GEF) that is capable of activating Rho A,
NGEF_HUMAN.  Background: EPHEXIN is a 710 amino acid protein that localizes to both the membrane and the cytoplasm and contains one SH3 domain, one PH domain and one DH domain. Expressed at high levels in brain and present at lower levels in lung tissue, EPHEXIN interacts with EphA4 and functions as a guanine nucleotide exchange factor (GEF) that is capable of activating Rho A,
Background: EPHEXIN is a 710 amino acid protein that localizes to both the membrane and the cytoplasm and contains one SH3 domain, one PH domain and one DH domain. Expressed at high levels in brain and present at lower levels in lung tissue, EPHEXIN interacts with EphA4 and functions as a guanine nucleotide exchange factor (GEF) that is capable of activating Rho A,
cytoplasm and contains one SH3 domain, one PH domain and one DH domain. Expressed at high levels in brain and present at lower levels in lung tissue, EPHEXIN interacts with EphA4 and functions as a guanine nucleotide exchange factor (GEF) that is capable of activating Rho A,
high levels in brain and present at lower levels in lung tissue, EPHEXIN interacts with EphA4 and functions as a guanine nucleotide exchange factor (GEF) that is capable of activating Rho A,
functions as a guanine nucleotide exchange factor (GEF) that is capable of activating Rho A,
Dog 1 and Cdo 12 and in thought to play a role in even guidenes and growth some colleges
Rac 1 and Cdc42 and is thought to play a role in axon guidance and growth cone collapse.
EPHEXIN is subject to Src-dependent phosphorylation, an event that increases the GEF activity
of EPHEXIN toward Rho A. Human EPHEXIN, which exists as multiple alternatively spliced
isoforms, shares a high degree of sequence homology with its mouse counterpart, suggesting
a conserved role between species.
25791
Neurotrophin Signaling Pathway
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200
For Research Use only
Liquid
1 μg/μL
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
ProClin
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
handled by trained staff only.
-20 °C

## Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months