# antibodies -online.com





## anti-GNG5 antibody (AA 21-65) (AbBy Fluor® 350)



Go to Product page

( )	1/0	r\ / I	014	
( )	ve	I V I	-v	V

Quantity:	100 μL	
Target:	GNG5	
Binding Specificity:	AA 21-65	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This GNG5 antibody is conjugated to AbBy Fluor® 350	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human GNG5	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Human,Mouse,Cow,Chicken,Rabbit	
Purification:	Purified by Protein A.	

#### Target Details

Target:	GNG5
Alternative Name:	GNG5 (GNG5 Products)

#### **Target Details**

Background:	Synonyms: FLJ92393, GBG5_HUMAN, Gng5, Guanine nucleotide-binding protein GI/GS/GO
	subunit gamma-5, OTTHUMP00000011474, OTTHUMP00000011565.
	Background: Heterotrimeric G proteins function to relay information from cell surface receptors
	to intracellular effectors. Each of a very broad range of receptors specifically detects an
	extracellular stimulus (i.e. a photon, pheromone, odorant, hormone or neurotransmitter), while
	the effectors (e.g. adenyl cyclase), which act to generate one or more intracellular messengers,
	are less numerous. In mammals, G protein alhfa, beta and gamma polypeptides are encoded by
	at least 16, 4 and 7 genes, respectively. Most interest in G proteins has been focused on their a
	subunits, since these proteins bind and hydrolyze GTP and most obviously regulate the activity
	of the best studied effectors. Evidence, however, has established an important regulatory role
	for the beta gamma subunits. It is becoming increasingly clear that different G protein
	complexes expressed in different tissues carry structurally distinct members of the gamma as
	well as the alhfa and beta subunits, and that preferential associations between members of
	subunit families increase G protein functional diversity.
Gene ID:	2787
Pathways:	Myometrial Relaxation and Contraction, SARS-CoV-2 Protein Interactome
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C

### Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months