antibodies -online.com





anti-IFT122 antibody (AA 1-100) (AbBy Fluor® 350)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL
Target:	IFT122
Binding Specificity:	AA 1-100
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IFT122 antibody is conjugated to AbBy Fluor® 350
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human IFT122
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Horse,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	IFT122
Alternative Name:	Ift122 (IFT122 Products)
Background:	Synonyms: CED, 122_HUMAN, T122, Intraflagellar transport 122 homolog Chlamydomonas,

Intraflagellar transport protein 122 homolog, SPG, WD repeat domain 10, WD repeat-containing protein 10, WD repeat-containing protein 140, WDR10, WDR10p, WDR140.

Background: IFT122 is a member of the WD repeat protein family. WD repeats are minimally conserved regions of approximately 40 amino acids typically bracketed by gly-his and trp-asp (GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction, apoptosis, and gene regulation. This cytoplasmic protein contains seven WD repeats and an AF-2 domain which function by recruiting coregulatory molecules and in transcriptional activation. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.

Pathways:

Tube Formation, Embryonic Body Morphogenesis

Application Details

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200 IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	