# antibodies .- online.com





## anti-ACSL4 antibody (AbBy Fluor® 350)



Go to Product page

$\sim$			
	N/P	r\/I	i⊢₩

Quantity:	100 μL
Target:	ACSL4
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ACSL4 antibody is conjugated to AbBy Fluor® 350
Application:	Western Blotting (WB)

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human FACL4/ACSL4
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Cow, Sheep, Pig, Horse, Chicken, Rabbit
Purification:	Purified by Protein A.

### **Target Details**

Target:	ACSL4
Alternative Name:	FacI4/AcsI4 (ACSL4 Products)
Background:	Synonyms: ACS 4, ACS4, ACSL 4, Acsl4, ACSL4_HUMAN, acyl CoA synthetase 4, Acyl CoA
	synthetase long chain family member 4, FACL 4, FACL4, Fatty acid Coenzyme A ligase, fatty
	acid Coenzyme A ligase long-chain 4, LACS 4, LACS4, Lignoceroyl CoA synthase, Long chain 4,
	long chain acyl CoA synthetase 4, long chain fatty acid CoA ligase 4, long chain fatty acid

Coenzyme A ligase 4, Long-chain acyl-CoA synthetase 4, Long-chain-fatty-acid--CoA ligase 4, MRX63, MRX68.

Background: Acyl-CoA synthetases, also known as long-chain fatty-acid CoA synthases (FACL) or palmitoyl-CoA ligases, include ACSL1-6, which are all single-pass membrane proteins localizing to the mitochondrion, microsome or peroxisome. FACL proteins are important for synthesis of cellular lipids and for -oxidation degradation. Specifically, ACSL proteins catalyze the activation of long-chain fatty acids to acyl-CoAs, which can be metabolized to form CO2, triacylglycerol (TAG), phospholipids (PL) and cholesteryl esters (CE). ACSL3 preferentially utilizes laurate, myristate, arachidonate and eicosapentaenoate among saturated and unsaturated long chain fatty acids. FACL3 is expressed as two isoforms in various tissues, including brain, heart, placenta, prostate, skeletal muscle, testis and thymus. FACL4 preferentially utilizes arachidonate and is abundant in steroidogenic tissues. FACL4 may modulate female fertility and uterine prostaglandin production.

Gene ID:

2182

#### Application Details

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Concentration:  $1 \mu g/\mu L$ Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. -20 °C Storage: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. Storage Comment: **Expiry Date:** 12 months