# antibodies -online.com





# anti-C6orf168 antibody (AA 221-320) (Alexa Fluor 350)



Go to Product page

$\sim$			
	N/P	r\/	i⊢₩

Quantity:	100 μL	
Target:	C6orf168 (FAXC)	
Binding Specificity:	AA 221-320	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This C6orf168 antibody is conjugated to Alexa Fluor 350	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Western Blotting (WB)	

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human C6orf168	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse	
Purification:	Purified by Protein A.	

# **Target Details**

Target:	C6orf168 (FAXC)	
Alternative Name:	C6orf168 (FAXC Products)	
Background:	kground: Synonyms: C6orf168, CF168_HUMAN, Chromosome 6 open reading frame 168, dJ273F20	

Uncharacterized protein C6orf168.

Background: Making up nearly 6 % of the human genome, chromosome 6 contains around 1,200 genes within 170 million base pairs of sequence. Deletion of a portion of the q arm of chromosome 6 is associated with early onset intestinal cancer suggesting the presence of a cancer susceptibility locus. Porphyria cutanea tarda is associated with chromosome 6 through the HFE gene which, when mutated, predisposes an individual to developing this porphyria. Notably, the PARK2 gene, which is associated with Parkinson's disease, and the genes encoding the major histocompatiblity complex proteins, which are key molecular components of the immune system and determine predisposition to rheumatic diseases, are also located on chromosome 6. Stickler syndrome, 21-hydroxylase deficiency and maple syrup urine disease are also associated with genes on chromosome 6. A bipolar disorder susceptibility locus has been identified on the q arm of chromosome 6. The C6orf168 gene product has been provisionally designated C6orf168 pending further characterization.

Gene ID:

84553

### **Application Details**

Application Notes:	

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months