# antibodies -online.com





# anti-GNAI1 antibody (AA 21-120) (Alexa Fluor 350)



Go to Product page

| $\sim$ |     |      |                               |
|--------|-----|------|-------------------------------|
|        | N/P | r\/I | $\Theta \backslash \Lambda /$ |

| Quantity:            | 100 μL  |
|----------------------|---|
| Target:              | GNAI1   |
| Binding Specificity: | AA 21-120   |
| Reactivity:          | Human   |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This GNAI1 antibody is conjugated to Alexa Fluor 350  |
| Application:         | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human G protein alpha inhibitor 1 |  |
|-----------------------|---|--|
| Isotype:              | IgG   |  |
| Predicted Reactivity: | Human,Mouse,Rat,Cow,Sheep,Pig,Zebrafish   |  |
| Purification:         | Purified by Protein A.  |  |

## **Target Details**

| Target:           | GNAI1   |  |
|-------------------|---|--|
| Alternative Name: | G protein alpha inhibitor 1 (GNAI1 Products)  |  |
| Background:       | Synonyms: Adenylate cyclase inhibiting G alpha protein, Adenylate cyclase inhibitory protein, |  |

| Adenylate cyclase-inhibiting G alpha protein, G protein alpha inhibiting 1, G protein alpha        |
|--|
| inhibiting activity polypeptide 1, Gi, Gi inhibitory G protein, Gi1 protein alpha subunit, GNAI 1, |
| GNAI1, GNAI1_HUMAN, Guanine nucleotide binding protein G protein alpha inhibiting activity         |
| polypeptide 1, Guanine nucleotide binding protein alpha inhibiting activity polypeptide 1,         |
| Guanine nucleotide binding protein Gi alpha 1 subunit, Guanine nucleotide-binding protein Gi       |
| subunit alpha-1, OTTHUMP00000161319, OTTHUMP00000208047.   |
|  |

Background: Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems. The G(i) proteins are involved in hormonal regulation of adenylate cyclase: they inhibit the cyclase in response to beta-adrenergic stimuli. The inactive GDP-bound form prevents the association of RGS14 with centrosomes and is required for the translocation of RGS14 from the cytoplasm to the plasma membrane. May play a role in cell division.

| Gene ID: | 2770 |
|----------|------|
|----------|------|

Pathways: G-protein mediated Events

## **Application Details**

| Application Notes: | IF(IHC-P) 1:50-200 |
|--------------------|--------------------|
|--------------------|--------------------|

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

#### Handling

| Liquid   |
|--|
| 1 μg/μL  |
| Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| ProClin  |
| This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| -20 °C   |
| Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
|  |

|   | l l  | 1:     |
|---|------|--------|
| - | เวทก | 111111 |
|   | land | 11110  |
|   |      |        |

Expiry Date:

12 months