# antibodies -online.com







# anti-GNPDA1 antibody (AA 201-289) (Alexa Fluor 350)



$\sim$			
	N/P	r\/I	i⊢₩

Quantity:	100 μL	
Target:	GNPDA1	
Binding Specificity:	AA 201-289	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This GNPDA1 antibody is conjugated to Alexa Fluor 350	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human GNPDA1
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Cow,Sheep,Pig
Purification:	Purified by Protein A.

## **Target Details**

Target:	GNPDA1	
Alternative Name:	GNPDA1 (GNPDA1 Products)	

## **Target Details**

Synonyms: EC 3.5.99.6, GlcN6P deaminase 1, Glucosamine 6 phosphate deaminase 1,	
Glucosamine 6 phosphate isomerase 1, Glucosamine-6-phosphate deaminase 1, Glucosamine-	
6-phosphate isomerase 1, GNPDA 1, GNPDA1, GNPI, GNPI1_HUMAN, HLN, KIAA0060, Oscillin,	
Putative glucosamine 6 phosphate isomerase.	
Background: During fertilization in mammals, the sperm activates the egg by causing an	
increase in the level of free cytoplasmic calcium concentration. This increased calcium	
concentration induces a characteristic series of oscillations that trigger egg activation and early	
embryo development. A hamster protein named oscillin is thought to be involved in this	
pathway. The enzyme glucosamine-6-phosphate isomerase (GNPI) or deaminase (GNPDA1)	
and the related protein GNPDA2 are the human homologs of hamster oscillin. GNPDA1 and	
GNPDA2 catalyze the conversion of GNP to fructose-6-phosphate and ammonia. Both proteins	
exist as homohexamers and are ubiquitously expressed with highest expression in testis, ovary	
and heart. Three isoforms of GNPDA2 are expressed due to alternative splicing events.	
10007	
IF(IHC-P) 1:50-200	
IF(IHC-F) 1:50-200	
IF(ICC) 1:50-200	
For Research Use only	
Liquid	
1 μg/μL	
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
50 % Glycerol.	
ProClin	
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
handled by trained staff only.	
-20 °C	
Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
12 months	