# antibodies -online.com







## anti-SKAP1 antibody (AA 101-200) (AbBy Fluor® 350)



$\sim$	
( )\/△	rview
$\cup$	1 410 44

Quantity:	100 μL	
Target:	SKAP1	
Binding Specificity:	AA 101-200	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This SKAP1 antibody is conjugated to AbBy Fluor® 350	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human SKAP55	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Human,Mouse,Dog,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	

### **Target Details**

Target:	SKAP1
Alternative Name:	Skap55 (SKAP1 Products)

#### **Target Details**

#### Background:

Synonyms: pp55, SCAP 1, SCAP 1, SKAP 1, SKAP 55, SKAP-55, Skap1, SKAP1\_HUMAN, SKAP55 adaptor protein, SRC family associated phosphoprotein 1, Src kinase associated phosphoprotein 1, Src kinase associated phosphoprotein 1, SRC kinase associated phosphoprotein of 55 kDa, Src kinase-associated phosphoprotein of 55 kDa, Src kinase-associated phosphoprotein of 55 kDa.

Background: Fyb (Fyn binding protein) and the anchoring proteins SKAP55 and SKAP55-R (SKAP55-related protein) associate with the tyrosine kinase p59fyn. SKAP55 and SKAP55-R bind to Fyb through their SH3 domains and function as substrates for p59Fyn in resting T cells. SKAP55 contains an N-terminal pleckstrin homology domain and a C-terminal SH3 domain binding motif of adjacent arginine and lysine residues followed by tandem tyrosines. SKAP55-R, similar in overall structure to SKAP55, contains a coiled-coil N-terminal domain. SKAP55 associates with SLAP-130, another component of the Fyn complex, which plays a role in the regulation of signaling events initiated by lymphocyte antigen receptors leading up to T cell activation. The human SKAP55 gene maps to chromosome 17q21.32 and encodes a 359 amino acid protein.

#### **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

$\vdash$	land	lına
	iaria	шц

Expiry Date:

12 months